

Optimal control of the bidomain system (III):
Existence of minimizers and first-order optimality conditions.
Revised version

Karl Kunisch and Marcus Wagner

1. Introduction.

In this work, we continue our investigations of optimal control problems for the bidomain system. After the study of the monodomain approximation of the equations and a thorough stability and regularity analysis of weak solutions for the full bidomain equations, as contained in the previous papers [KUNISCH/WAGNER 12] and [KUNISCH/WAGNER 11], we are now in position to analyze the related control problems with respect to the existence of minimizers as well as to provide a rigorous proof of the first-order necessary optimality conditions.

Let $\Omega \subset \mathbb{R}^3$ be a bounded domain and $T > 0$ a fixed time horizon. Then the bidomain system, representing a well-accepted description of the electrical activity of the heart, is given by⁰¹⁾

$$\frac{\partial \Phi_{tr}}{\partial t} + I_{ion}(\Phi_{tr}, W) - \operatorname{div}(M_i \nabla \Phi_i) = I_i \quad \text{for almost all } (x, t) \in \Omega \times [0, T]; \quad (1.1)$$

$$\frac{\partial \Phi_{tr}}{\partial t} + I_{ion}(\Phi_{tr}, W) + \operatorname{div}(M_e \nabla \Phi_e) = -I_e \quad \text{for almost all } (x, t) \in \Omega \times [0, T]; \quad (1.2)$$

$$\frac{\partial W}{\partial t} + G(\Phi_{tr}, W) = 0 \quad \text{for almost all } (x, t) \in \Omega \times [0, T]; \quad (1.3)$$

$$\mathbf{n}^T M_i \nabla \Phi_i = 0 \quad \text{for all } (x, t) \in \partial\Omega \times [0, T]; \quad (1.4)$$

$$\mathbf{n}^T M_e \nabla \Phi_e = 0 \quad \text{for all } (x, t) \in \partial\Omega \times [0, T]; \quad (1.5)$$

$$\Phi_{tr}(x, 0) = \Phi_i(x, 0) - \Phi_e(x, 0) = \Phi_0(x) \quad \text{and} \quad W(x, 0) = W_0(x) \quad \text{for almost all } x \in \Omega. \quad (1.6)$$

In this model, Ω represents the spatial domain occupied by the cardiac muscle, the variables Φ_i and Φ_e denote the intracellular and extracellular electric potentials, and $\Phi_{tr} = \Phi_i - \Phi_e$ is the transmembrane potential. The anisotropic electric properties of the intracellular and the extracellular tissue parts are modeled by conductivity tensors M_i and M_e . The specification of the model for the ionic current I_{ion} in (1.1) and (1.2) and the gating function G in (1.3) will be made below. We shall consider three so-called two-variable models wherein I_{ion} and G depend on Φ_{tr} as well as on a single gating variable W , which describes in a cumulative way the effects of the ion transport through the cell membranes (see Subsection 2.2.). Finally, the inhomogeneities I_i and I_e represent the intracellular and extracellular stimulation currents, respectively. We shall investigate optimal control problems of the form

$$(P) \quad F(\Phi_{tr}, \Phi_e, W, I_e) = \int_0^T \int_{\Omega} r(x, t, \Phi_{tr}(x, t), \Phi_e(x, t), W(x, t)) \, dx \, dt + \frac{\mu}{2} \int_0^T \int_{\Omega_{con}} I_e(x, t)^2 \, dx \, dt \longrightarrow \inf! \quad (1.7)$$

subject to the bidomain equations (1.1) – (1.6) in its weak formulation (see (2.1) – (2.4) below)

$$\text{and the control restriction } I_e \in \mathcal{C} \quad (1.8)$$

⁰¹⁾ The bidomain model has been considered first in [TUNG 78]. A detailed introduction may be found e. g. in [SUNDNES/LINES/CAI/NIELSEN/MARDAL/TVEITO 06], pp. 21 – 56.

where Ω_{con} is a Lipschitz subdomain of Ω and

$$\mathcal{C} = \{ QI \mid I \in L^\infty[(0, T), L^2(\Omega)], \text{supp}(I) \subseteq \Omega_{con} \times [0, T], \quad (1.9)$$

$$|I(x, t)| \leq R \ (\forall) (x, t) \in \Omega_T \} \subset L^\infty[(0, T), L^2(\Omega)].$$

For the description of the control domain, the linear operator $Q: L^2(\Omega_T) \rightarrow L^2(\Omega_T)$ defined by

$$QI(x, t) = I(x, t) - \mathbb{1}_{\Omega_{con}}(x) \cdot \frac{1}{|\Omega_{con}|} \int_{\Omega_{con}} I(\tilde{x}, t) d\tilde{x} \quad (1.10)$$

has been used. When applied to a function I with $\text{supp}(I) \subseteq \Omega_{con} \times [0, T]$, Q extends by zero the orthogonal projection onto the complement of the subspace $\{Z \mid \int_{\Omega_{con}} Z(\tilde{x}, t) d\tilde{x} = 0 \text{ for a. a. } t \in (0, T)\} \subset L^2[(0, T), L^2(\Omega_{con})]$. Consequently, for $I_e \in \mathcal{C}$, we have

$$\int_{\Omega} I_e(x, t) dx = \int_{\Omega_{con}} I_e(x, t) dx = 0 \quad \text{for almost all } t \in (0, T), \quad (1.11)$$

what guarantees the solvability of the state equations (cf. Theorem 2.3. below). In problem (P), the extracellular excitation I_e acts as control, which is allowed to be applied on the subdomain Ω_{con} only.⁰²⁾ The pointwise constraint within the description (1.9) of \mathcal{C} is included due to the obvious fact that one cannot apply arbitrary large electrical stimulations to living tissue without damaging it. In mathematical terms, this restriction is necessary in order to establish a stability estimate for the bidomain system (Theorem 2.4.).

Due to the complex dynamical behaviour of the state equations, an appropriate choice of the integrand r within the first term of the objective (1.7) for concrete applications is quite delicate. With arrhythmia or tachycardia in mind, it could be chosen as $r(x, t, \varphi, \eta, w) = (\varphi - \Phi_{des}(t))^2$ where Φ_{des} denotes some desired trajectory for the controlled state Φ_{tr} , which is part of a solution of (1.1) – (1.5) as well, cf. [NAGAI/IAH/KUNISCH/PLANK 11B]. The second term expresses the requirement that — regardless of whether the pointwise restriction within (1.9) is active — the overall stimulus should be as small as possible. Consequently, solutions with little intervention to the cardiac system are favored.

Besides an existence theorem for global minimizers (Theorem 3.4.), the main result of the present paper is the rigorous proof of the following *set of first-order necessary optimality conditions* for sufficiently regular local minimizers $(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e)$ of (P), consisting of the variational inequality

$$\int_0^T \int_{\Omega_{con}} (\mu \hat{I}_e - Q P_2) \cdot (I_e - \hat{I}_e) dx dt \geq 0 \quad \text{for all admissible controls } I_e \quad (1.12)$$

and the adjoint system⁰³⁾

$$\int_0^T \int_{\Omega} \left(-\frac{\partial P_1}{\partial t} + \frac{\partial I_{ion}}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{W}) P_1 + \frac{\partial G}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{W}) P_3 \right) \psi dx dt + \int_0^T \int_{\Omega} \nabla \psi^T M_i (\nabla P_1 + \nabla P_2) dx dt \quad (1.13)$$

$$= - \int_0^T \int_{\Omega} \left(\frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right) \psi dx dt \quad \forall \psi \in L^2[(0, T), W^{1,2}(\Omega)], \quad P_1(x, T) \equiv 0;$$

$$\int_0^T \int_{\Omega} \nabla \psi^T M_i \nabla P_1 dx dt + \int_0^T \int_{\Omega} \nabla \psi^T (M_i + M_e) \nabla P_2 dx dt = - \int_0^T \int_{\Omega} \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \psi dx dt \quad (1.14)$$

$$\forall \psi \in L^2[(0, T), W^{1,2}(\Omega)] \text{ with } \int_{\Omega} \psi(x, t) dx = 0 \text{ for a. a. } t \in (0, T), \quad \int_{\Omega} P_2(x, t) dx = 0 \ (\forall) t \in (0, T);$$

$$\int_0^T \int_{\Omega} \left(-\frac{\partial P_3}{\partial t} + \frac{\partial I_{ion}}{\partial w}(\hat{\Phi}_{tr}, \hat{W}) P_1 + \frac{\partial G}{\partial w}(\hat{\Phi}_{tr}, \hat{W}) P_3 \right) \psi dx dt = - \int_0^T \int_{\Omega} \left(\frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right) \psi dx dt \quad (1.15)$$

$$\forall \psi \in L^2[(0, T), L^2(\Omega)], \quad P_3(x, T) \equiv 0$$

⁰²⁾ For physiological reasons, the intracellular excitation I_i must be set zero.

⁰³⁾ Within the functions $r(x, t, \varphi, \eta, w)$, $I_{ion}(\varphi, w)$ and $G(\varphi, w)$, the real variables φ , η and w are the placeholders for Φ_{tr} , Φ_e and W , respectively.

for the multipliers P_1 , P_2 and P_3 related to the weak state equations (2.1), (2.2) and (2.3) below, respectively (Theorem 5.2.). The proof, which will be given by fitting the problem (P) into the framework of *weakly singular problems* in the sense of Ito/Kunisch (see [ITO/KUNISCH 08], p. 17 f.), is based on two main ingredients. The first one is a stability estimate for the primal equations (Theorem 2.4.), whose proof has been already provided in the previous publication [KUNISCH/WAGNER 11]. Secondly, we need an existence proof for weak solutions of the adjoint system, which is contained in the present paper (Theorem 4.2.). In difference to the monodomain approximation considered in [KUNISCH/WAGNER 12], the proof of the optimality conditions requires additional regularity of the minimizer $(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e)$ in the case of the full bidomain system.

In the literature, only a few studies related to the optimal control of the bidomain system are available as yet, mostly restricted to the monodomain approximation. We mention [AINSEBA/BENDAHMANE/RUIZ-BAIER 12], [BRANDAÕ/FERNÁNDEZ-CARA/MAGALHÃES/ROJAS-MEDAR 08], [KUNISCH/NAGAIAH/WAGNER 12], [MUZDEKA/BARBIERI 05], [NAGAIAH/KUNISCH 11], [NAGAIAH/KUNISCH/PLANK 11A] and [NAGAIAH/KUNISCH/PLANK 11B] and refer to [KUNISCH/WAGNER 12], p. 1527, for a closer discussion. Numerical work concerning open-loop control of the bidomain equations with the goal of dampening of excitation and reentry waves has been realized in [KUNISCH/NAGAIAH/WAGNER 12], [NAGAIAH/KUNISCH 11], [NAGAIAH/KUNISCH/PLANK 11A] and [NAGAIAH/KUNISCH/PLANK 11B]. The problems were treated with gradient and Newton-type techniques applied to FEM discretizations of the state equations.

The paper is structured in the following way. In Section 2, the solution concepts for the bidomain equations are outlined. We present the ionic models to be used and summarize the existence and stability theorems for weak solutions of (1.1) – (1.6). Then, in Section 3, we restate the optimal control problem (1.7) – (1.8) within function spaces, subsequently analyzing the structure of the feasible domain and establishing the existence of global minimizers. Section 4 is concerned with the derivation of the adjoint system and the existence proof for a weak solution of it. Finally, in Section 5, we state and prove the first-order necessary optimality conditions for the control problem.

Notations.

We denote by $L^p(\Omega)$ the space of functions, which are in the p th power integrable ($1 \leq p < \infty$), or are measurable and essentially bounded ($p = \infty$), and by $W^{1,p}(\Omega)$ the Sobolev space of functions $\psi: \Omega \rightarrow \mathbb{R}$ which, together with their first-order weak partial derivatives, belong to the space $L^p(\Omega, \mathbb{R})$ ($1 \leq p < \infty$). For spaces of Bochner integrable mappings, e. g. $L^2[(0, T), W^{1,2}(\Omega)]$, we refer to the summary in [KUNISCH/WAGNER 12], p. 1542. Ω_T is an abbreviation for $\Omega \times [0, T]$. The gradient ∇ is always taken only with respect to the spatial variables x . The characteristic function of the set $A \subseteq \mathbb{R}^3$ is defined as $\mathbf{1}_A: \mathbb{R}^3 \rightarrow \mathbb{R}$ with $\mathbf{1}_A(x) = 1 \iff x \in A$ and $\mathbf{1}_A(x) = 0 \iff x \notin A$. Finally, the nonstandard abbreviation “ $(\forall) t \in A$ ” has to be read as “for almost all $t \in A$ ” or “for all $t \in A$ except for a Lebesgue null set”, and the symbol \mathfrak{o} denotes, depending on the context, the zero element or the zero function of the underlying space.

2. Weak solutions of the bidomain system.

2.1. Parabolic-elliptic form of the bidomain system; strong and weak solutions.

It is well-known that the bidomain system (1.1) – (1.6) can be equivalently stated in parabolic-elliptic form, cf. [BOURGAULT/COUDIÈRE/PIERRE 09], p. 459, and [KUNISCH/WAGNER 11], p. 4, (2.1) – (2.9). In its weak formulation, the system reads as follows:

$$\int_{\Omega} \left(\frac{\partial \Phi_{tr}}{\partial t} \cdot \psi + \nabla \psi^T M_i (\nabla \Phi_{tr} + \nabla \Phi_e) + I_{ion}(\Phi_{tr}, W) \psi \right) dx = \int_{\Omega} I_i \psi dx \quad (2.1)$$

$$\forall \psi \in W^{1,2}(\Omega), \text{ for a. a. } t \in (0, T);$$

$$\int_{\Omega} \left(\nabla \psi^T M_i \nabla \Phi_{tr} + \nabla \psi^T (M_i + M_e) \nabla \Phi_e \right) dx = \int_{\Omega} (I_i + I_e) \psi dx \quad (2.2)$$

$$\forall \psi \in W^{1,2}(\Omega) \text{ with } \int_{\Omega} \psi(x) dx = 0, \text{ for a. a. } t \in (0, T);$$

$$\int_{\Omega} \left(\frac{\partial W}{\partial t} + G(\Phi_{tr}, W) \right) \psi dx = 0 \quad \forall \psi \in L^2(\Omega), \text{ for a. a. } t \in (0, T); \quad (2.3)$$

$$\Phi_{tr}(x, 0) = \Phi_0(x) \quad \text{and} \quad W(x, 0) = W_0(x) \quad \text{for almost all } x \in \Omega. \quad (2.4)$$

Throughout the paper, the following assumptions about the data will be made:

Assumptions 2.1. (Basic assumptions on the data)

1) $\Omega \subset \mathbb{R}^3$ is a bounded Lipschitz domain.

2) $M_i, M_e : \text{cl}(\Omega) \rightarrow \mathbb{R}^{3 \times 3}$ are symmetric, positive definite matrix functions with $L^\infty(\Omega)$ -coefficients, obeying uniform ellipticity conditions:

$$0 \leq \mu_1 \|\xi\|^2 \leq \xi^T M_i(x) \xi \leq \mu_2 \|\xi\|^2 \quad \text{and} \quad 0 \leq \mu_1 \|\xi\|^2 \leq \xi^T M_e(x) \xi \leq \mu_2 \|\xi\|^2 \quad \forall \xi \in \mathbb{R}^3 \quad \forall x \in \Omega \quad (2.5)$$

with $\mu_1, \mu_2 > 0$.

The notions of strong and weak solutions are as follows:

Definition 2.2. 1)⁰⁴ **(Strong solution of the bidomain system)** A triple (Φ_{tr}, Φ_e, W) is called a strong solution of the bidomain system (2.1) – (2.4) on $[0, T]$ iff the functions Φ_{tr}, Φ_e and W satisfy (2.1) – (2.4) and belong to the spaces

$$\Phi_{tr} \in L^2[(0, T), W^{2,2}(\Omega)] \cap W^{1,2}[(0, T), L^2(\Omega)]; \quad (2.6)$$

$$\Phi_e \in L^2[(0, T), W^{2,2}(\Omega)]; \quad (2.7)$$

$$W \in W^{1,2}[(0, T), L^2(\Omega)] \quad (2.8)$$

where $\int_{\Omega} \Phi_e(x, t) dx = 0$ holds for almost all $t \in (0, T)$.

2)⁰⁵ **(Weak solution of the bidomain system)** A triple (Φ_{tr}, Φ_e, W) is called a weak solution of the bidomain system (2.1) – (2.4) on $[0, T]$ iff the functions Φ_{tr}, Φ_e and W satisfy (2.1) – (2.4) and belong to the spaces

$$\Phi_{tr} \in C^0[[0, T], L^2(\Omega)] \cap L^2[(0, T), W^{1,2}(\Omega)] \cap L^4(\Omega_T); \quad (2.9)$$

$$\Phi_e \in L^2[(0, T), W^{1,2}(\Omega)]; \quad (2.10)$$

$$W \in C^0[[0, T], L^2(\Omega)] \quad (2.11)$$

where $\int_{\Omega} \Phi_e(x, t) dx = 0$ holds for almost all $t \in (0, T)$.

2.2. Two-variable models for the ionic current.

For the ionic current I_{ion} and the function G within the gating equation, the following three models will be considered:

⁰⁴) Slightly modified from [BOURGAULT/COUDIÈRE/PIERRE 09], p. 469, Definition 18.

⁰⁵) [BOURGAULT/COUDIÈRE/PIERRE 09], p. 472, Definition 26.

a) *The Rogers-McCulloch model.*⁰⁶⁾

$$I_{ion}(\varphi, w) = b \cdot \varphi (\varphi - a) (\varphi - 1) + \varphi \cdot w = b \varphi^3 - (a + 1) b \varphi^2 + a b \varphi + \varphi w; \quad (2.12)$$

$$G(\varphi, w) = \varepsilon w - \varepsilon \kappa \varphi \quad (2.13)$$

with $0 < a < 1$, $b > 0$, $\kappa > 0$ and $\varepsilon > 0$. Consequently, the gating variable obeys the linear ODE

$$\partial W / \partial t + \varepsilon W = \varepsilon \kappa \Phi_{tr}. \quad (2.14)$$

b) *The FitzHugh-Nagumo model.*⁰⁷⁾

$$I_{ion}(\varphi, w) = \varphi (\varphi - a) (\varphi - 1) + w = \varphi^3 - (a + 1) \varphi^2 + a \varphi + w; \quad (2.15)$$

$$G(\varphi, w) = \varepsilon w - \varepsilon \kappa \varphi \quad (2.16)$$

with $0 < a < 1$, $\kappa > 0$ and $\varepsilon > 0$. Consequently, the gating variable obeys the same linear ODE (2.14) as before.

c) *The linearized Aliev-Panfilov model.*⁰⁸⁾

$$I_{ion}(\varphi, w) = b \cdot \varphi (\varphi - a) (\varphi - 1) + \varphi \cdot w = b \varphi^3 - (a + 1) b \varphi^2 + a b \varphi + \varphi w; \quad (2.17)$$

$$G(\varphi, w) = \varepsilon w - \varepsilon \kappa ((a + 1) \varphi - \varphi^2) \quad (2.18)$$

with $0 < a < 1$, $b > 0$, $\kappa > 0$ and $\varepsilon > 0$. The linear ODE for the gating variable is

$$\partial W / \partial t + \varepsilon W = \varepsilon \kappa ((a + 1) \Phi_{tr} - \Phi_{tr}^2). \quad (2.19)$$

2.3. Existence and uniqueness of weak solutions; the stability estimate.

In [KUNISCH/WAGNER 11], the following results about weak solutions of the bidomain system (2.1) – (2.4) have been obtained:

Theorem 2.3. (Existence and uniqueness of weak solutions)⁰⁹⁾ *Assume that the data within (2.1) – (2.4) obey Assumptions 2.1., and specify the Rogers-McCulloch or the FitzHugh-Nagumo model. Then the bidomain system (2.1) – (2.4) admits for arbitrary initial values $\Phi_0 \in L^2(\Omega)$, $W_0 \in L^4(\Omega)$ and inhomogeneities $I_i, I_e \in L^\infty[(0, T), (W^{1,2}(\Omega))^*]$, which satisfy the compatibility condition*

$$\int_{\Omega} \left(I_i(x, t) + I_e(x, t) \right) dx = 0 \quad \text{for almost all } t \in (0, T), \quad (2.20)$$

a uniquely determined weak solution (Φ_{tr}, Φ_e, W) on $[0, T]$ according to Definition 2.2., 2). If the linearized Aliev-Panfilov model is specified, this assertion remains true provided that W_0 belongs to $W^{1,3/2}(\Omega)$ instead of $L^4(\Omega)$.

⁰⁶⁾ Introduced in [ROGERS/MCCULLOCH 94].

⁰⁷⁾ See [FITZHUGH 61] together with [NAGUMO/ARIMOTO/YOSHIKAWA 62].

⁰⁸⁾ The model, which appears to be a linearization of the original model derived in [ALIEV/PANFILOV 96], is taken from [BOURGAULT/COUDIÈRE/PIERRE 09], p. 480.

⁰⁹⁾ [BOURGAULT/COUDIÈRE/PIERRE 09], p. 473, Theorem 30, together with [KUNISCH/WAGNER 11], p. 8, Theorem 2.8., slightly modified. An error within the proof of this and the next theorem will be fixed in a subsequent publication.

In fact, a closer regularity analysis reveals that, under the assumptions of Theorem 2.3., the components (Φ_{tr}, W) of a given weak solution of the bidomain system belong to $(L^2[(0, T), L^6(\Omega)] \cap L^q[(0, T), L^r(\Omega)]) \times C^0[[0, T], L^4(\Omega)]$ in the case of the Rogers-McCulloch or the FitzHugh-Nagumo model and to $(L^2[(0, T), L^6(\Omega)] \cap L^q[(0, T), L^r(\Omega)]) \times C^0[[0, T], L^{8/3}(\Omega)]$ in the case of the linearized Aliev-Panfilov model where $1 < q < \infty$ and $4 \leq r < 6$.

Theorem 2.4. (Stability estimate for weak solutions)¹⁰⁾ *Assume that the data within (2.1) – (2.4) obey Assumptions 2.1., and specify the Rogers-McCulloch or the FitzHugh-Nagumo model. Consider two weak solutions $(\Phi_{tr}', \Phi_e', W')$, $(\Phi_{tr}'', \Phi_e'', W'')$ of (2.1) – (2.4), which correspond to initial values $\Phi_0' = \Phi_0'' = \Phi_0 \in L^2(\Omega)$, $W_0' = W_0'' = W_0 \in L^4(\Omega)$ and inhomogeneities I_i', I_e', I_i'' and $I_e'' \in L^\infty[(0, T), (W^{1,2}(\Omega))^*]$ with*

$$\int_{\Omega} \left(I_i'(x, t) + I_e'(x, t) \right) dx = \int_{\Omega} \left(I_i''(x, t) + I_e''(x, t) \right) dx = 0 \quad \text{for almost all } t \in (0, T), \quad (2.21)$$

whose norms are bounded by $R > 0$. Then the following estimate holds:

$$\begin{aligned} & \| \Phi_{tr}' - \Phi_{tr}'' \|_{L^2[(0, T), W^{1,2}(\Omega)]}^2 + \| \Phi_{tr}' - \Phi_{tr}'' \|_{C^0[[0, T], L^2(\Omega)]}^2 \\ & + \| \Phi_{tr}' - \Phi_{tr}'' \|_{W^{1,4/3}[(0, T), (W^{1,2}(\Omega))^*]}^2 + \| \Phi_e' - \Phi_e'' \|_{L^2[(0, T), W^{1,2}(\Omega)]}^2 \\ & + \| W' - W'' \|_{L^2[(0, T), L^2(\Omega)]}^2 + \| W' - W'' \|_{C^0[[0, T], L^2(\Omega)]}^2 + \| W' - W'' \|_{W^{1,2}[(0, T), L^2(\Omega)]}^2 \\ & \leq C \left(\| I_i' - I_i'' \|_{L^\infty[(0, T), (W^{1,2}(\Omega))^*]}^2 + \| I_e' - I_e'' \|_{L^\infty[(0, T), (W^{1,2}(\Omega))^*]}^2 \right). \end{aligned} \quad (2.22)$$

The constant $C > 0$ does not depend on I_i', I_e', I_i'' and I_e'' but possibly on Ω, R, Φ_0 and W_0 . If the linearized Aliev-Panfilov model is specified then the assertion remains true provided that $W_0' = W_0'' = W_0$ belong to $W^{1,3/2}(\Omega)$ instead of $L^4(\Omega)$.

The assumptions in Theorems 2.3. – 2.4. are in accordance to the analytical framework wherein the control problem (P) will be studied in the next sections.

3. The optimal control problem.

3.1. Formulation of the problem within function spaces.

In order to provide a precise statement of the optimal control problem (1.7) – (1.8) within an appropriate function space framework, we introduce the following spaces:

$$X_1 = L^2[(0, T), W^{1,2}(\Omega)]; \quad X_2 = X_1 \cap \left\{ Z \mid \int_{\Omega} Z(x, t) dx = 0 \quad (\forall) t \in (0, T) \right\}; \quad (3.1)$$

$$X_3 = L^2[(0, T), L^2(\Omega)]; \quad X_4 = L^\infty[(0, T), L^2(\Omega)]. \quad (3.2)$$

We will further specify the subspaces

$$\tilde{X}_1 = X_1 \cap W^{1,4/3}[(0, T), (W^{1,2}(\Omega))^*] \cap C^0[[0, T], L^2(\Omega)]; \quad \tilde{X}_2 = X_2; \quad (3.3)$$

$$\tilde{X}_3 = X_3 \cap W^{1,2}[(0, T), (L^2(\Omega))^*] \cap C^0[[0, T], L^2(\Omega)], \quad (3.4)$$

¹⁰⁾ [KUNISCH/WAGNER 11], p. 7 f., Theorem 2.7., slightly modified.

which contain all polynomials and, consequently, lie dense in X_1 , X_2 and X_3 , as well as the target spaces

$$Z_1 = L^{4/3}[(0, T), (W^{1,2}(\Omega))^*]; \quad Z_2 = L^2[(0, T), (W^{1,2}(\Omega))^*]; \quad (3.5)$$

$$Z_3 = L^2[(0, T), (L^2(\Omega))^*]; \quad Z_4 = Z_5 = L^2(\Omega). \quad (3.6)$$

The quadruples $(\Phi_{tr}, \Phi_e, W, I_e)$ of state and control variables will be chosen from the space $\tilde{X}_1 \times \tilde{X}_2 \times \tilde{X}_3 \times X_4$. Recall the definition of $Q: L^2(\Omega_T) \rightarrow L^2(\Omega_T)$ as

$$QI(x, t) = I(x, t) - \mathbf{1}_{\Omega_{con}}(x) \cdot \frac{1}{|\Omega_{con}|} \int_{\Omega_{con}} I(\tilde{x}, t) d\tilde{x}. \quad (3.7)$$

With the aid of the operators

$$F: X_1 \times X_2 \times X_3 \times X_4 \rightarrow \mathbb{R}; \quad (3.8)$$

$$E_1: \tilde{X}_1 \times \tilde{X}_2 \times \tilde{X}_3 \rightarrow Z_1; \quad E_2: \tilde{X}_1 \times \tilde{X}_2 \times X_4 \rightarrow Z_2; \quad E_3: \tilde{X}_1 \times \tilde{X}_3 \rightarrow Z_3; \quad (3.9)$$

$$E_4: \tilde{X}_1 \rightarrow Z_4; \quad E_5: \tilde{X}_3 \rightarrow Z_5, \quad (3.10)$$

the problem (P) will be restated now in the following way:

$$(P) \quad F(\Phi_{tr}, \Phi_e, W, I_e) = \int_0^T \int_{\Omega} r(x, t, \Phi_{tr}(x, t), \Phi_e(x, t), W(x, t)) dx dt \quad (3.11)$$

$$+ \frac{\mu}{2} \cdot \int_0^T \int_{\Omega} I_e(x, t)^2 dx dt \rightarrow \inf!;$$

$$E_1(\Phi_{tr}, \Phi_e, W) = \mathbf{o} \iff \int_{\Omega} \left(\frac{\partial \Phi_{tr}}{\partial t} + I_{ion}(\Phi_{tr}, W) \right) \psi dx + \int_{\Omega} \nabla \psi^T M_i (\nabla \Phi_{tr} + \nabla \Phi_e) dx = 0 \quad (3.12)$$

$$\forall \psi \in W^{1,2}(\Omega) \quad (\forall) t \in (0, T);$$

$$E_2(\Phi_{tr}, \Phi_e, I_e) = \mathbf{o} \iff \int_{\Omega} \left(\nabla \psi^T M_i \nabla \Phi_{tr} + \nabla \psi^T (M_i + M_e) \nabla \Phi_e \right) dx - \int_{\Omega} I_e \psi dx = 0 \quad (3.13)$$

$$\forall \psi \in W^{1,2}(\Omega) \quad \text{with} \quad \int_{\Omega} \psi(x) dx = 0 \quad (\forall) t \in (0, T);$$

$$E_3(\Phi_{tr}, W) = \mathbf{o} \iff \int_{\Omega} \left(\frac{\partial W(t)}{\partial t} + G(\Phi_{tr}(t), W(t)) \right) \psi dx = 0 \quad \forall \psi \in L^2(\Omega) \quad (\forall) t \in (0, T); \quad (3.14)$$

$$E_4(\Phi_{tr}) = \mathbf{o} \iff \Phi_{tr}(x, 0) - \Phi_0(x) = 0 \quad (\forall) x \in \Omega; \quad (3.15)$$

$$E_5(W) = \mathbf{o} \iff W(x, 0) - W_0(x) = 0 \quad (\forall) x \in \Omega; \quad (3.16)$$

$$I_e \in \mathcal{C} = \{ QI \mid I \in L^\infty[(0, T), L^2(\Omega)], \text{supp}(I) \subseteq \Omega_{con} \times [0, T], \quad (3.17)$$

$$|I(x, t)| \leq R \quad (\forall) (x, t) \in \Omega_T \} \subset L^\infty[(0, T), L^2(\Omega)].$$

Assumptions 2.1. are imposed on the data of problem (P). The numbers $T > 0$, $\mu > 0$ and $R > 0$ as well as the Lipschitz subdomain $\Omega_{con} \subseteq \Omega$ are fixed. The functions I_{ion} and G will be specified according to any of the models from Subsection 2.2. In the case of the Rogers-McCulloch or the FitzHugh-Nagumo model, we fix initial values $\Phi_0 \in L^2(\Omega)$ and $W_0 \in L^4(\Omega)$ while in the case of the linearized Aliev-Panfilov model, $\Phi_0 \in L^2(\Omega)$ and $W_0 \in W^{1,3/2}(\Omega)$ will be used. Concerning the objective functional F , we assume the integrand

$$r(x, t, \varphi, \eta, w): \Omega \times [0, T] \times \mathbb{R}^3 \rightarrow \mathbb{R} \quad (3.18)$$

to be measurable with respect to x and t and continuous with respect to φ , η and w . With regard to (3.7) and (3.17), in the second term of F the original integration domain Ω_{con} from (1.7) can be replaced by Ω .

3.2. Structure of the feasible domain.

Proposition 3.1. *For the problem (3.11) – (3.17), the control-to-state-mapping $\mathcal{C} \ni I_e \mapsto (\Phi_{tr}, \Phi_e, W) \in X_1 \times X_2 \times X_3$ is well-defined.*

Proof. Recall that $\int_{\Omega} I_e(x, t) dx = 0$ for almost all $t \in (0, T)$. Consequently, the data within the problem (3.11) – (3.17) satisfy the assumptions of Theorem 2.3. with $I_i = \mathbf{o}$, and the existence of a uniquely determined weak solution (Φ_{tr}, Φ_e, W) of the bidomain system is guaranteed for any feasible control $I_e \in \mathcal{C} \subset L^\infty[(0, T), L^2(\Omega)]$. ■

Proposition 3.2. *The control domain $\mathcal{C} \subset L^\infty(\Omega_T)$ forms a closed, convex, weak*-sequentially compact subset of the space X_4 .*

Proof. Obviously, \mathcal{C} is a convex subset of X_4 . In order to confirm closedness, consider a norm-convergent sequence $\{Q I^N\}$ with members in $\mathcal{C} \cap X_4$ and limit element \hat{I} . Since the sequence $\{I^N\}$ of the generating functions is uniformly bounded in $L^\infty[(0, T), L^2(\Omega_{con})]$, it admits a weak*-convergent subsequence $I^{N'}$ with a limit element \tilde{I} still satisfying the conditions $\text{supp}(\tilde{I}) \subseteq \Omega_{con} \times [0, T]$ and $|\tilde{I}(x, t)| \leq R$ (\forall) $(x, t) \in \Omega_T$. The weak*-continuity of the operator Q implies then $Q I^{N'} \xrightarrow{*} X_4 Q \tilde{I}$ and $\hat{I} = Q \tilde{I} \in \mathcal{C}$. Now the weak*-sequential compactness of \mathcal{C} is obtained from [ROLEWICZ 76], p. 301, Theorem VI.6.6., together with p. 152, Theorem IV.4.11. Finally, $\|I\|_{L^\infty(\Omega_T)} \leq R$ implies $\|Q I\|_{L^\infty(\Omega_T)} \leq 2R$, and \mathcal{C} belongs even to $L^\infty(\Omega_T)$. ■

Proposition 3.3. *The feasible domain \mathcal{B} of the problem (3.11) – (3.17) is nonempty and closed with respect to the following topology in $X_1 \times X_2 \times X_3 \times X_4$: weak convergence with respect to the first three components, and weak*-convergence with respect to the fourth component.*

Proof. The existence of feasible solutions follows via Theorem 2.3. from Proposition 3.1. Consider now a sequence of feasible solutions $\{(\Phi_{tr}^N, \Phi_e^N, W^N, I_e^N)\}$ with $\Phi_{tr}^N \rightharpoonup X_1 \hat{\Phi}_{tr}$, $\Phi_e^N \rightharpoonup X_2 \hat{\Phi}_e$, $W^N \rightharpoonup X_3 \hat{W}$ and $I_e^N \xrightarrow{*} X_4 \hat{I}_e$. From Proposition 3.2. we already know that \hat{I}_e belongs to \mathcal{C} . Further, from [KUNISCH/WAGNER 11], p. 7, Theorem 2.6., we obtain uniform bounds with respect to N for the norms of Φ_{tr}^N , Φ_e^N , W^N , $\partial\Phi_{tr}^N/\partial t$ and $\partial W^N/\partial t$, implying weak convergence of $\partial\Phi_{tr}^{N'}/\partial t$, $\nabla\Phi_{tr}^{N'}$ and $\partial W^{N'}/\partial t$ as well as a. e. pointwise convergence of $\Phi_{tr}^{N'}$ on Ω_T along a suitable subsequence. Consequently, passing to the limit $N' \rightarrow \infty$ in (2.1) – (2.4), we may confirm that $(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W})$ solves the bidomain system with right-hand sides $I_i = \mathbf{o}$ and \hat{I}_e . ■

3.3. Existence of global minimizers.

Theorem 3.4. (Existence of global minimizers in (P)) *We impose the assumptions from Subsection 3.1. on the data of the problem (3.11) – (3.17). Assume further that the integrand $r(x, t, \varphi, \eta, w): \Omega \times [0, T] \times \mathbb{R}^3 \rightarrow \mathbb{R}$ is bounded from below and convex with respect to φ , η and w . Then the problem (3.11) – (3.17) admits a global minimizer.*

Proof. Since r is bounded from below, the problem (3.11) – (3.17) admits a minimizing sequence $\{(\Phi_{tr}^N, \Phi_e^N, W^N, I_e^N)\}$ of feasible solutions. Due to the uniform boundedness of $\|I_e^N\|_{X_4}$ with respect to N , the norms $\|\Phi_{tr}^N\|_{X_1}$, $\|\Phi_e^N\|_{X_2}$ and $\|W^N\|_{X_3}$ are uniformly bounded as well (cf. again [KUNISCH/WAGNER 11], p. 7, Theorem 2.6.), and we may pass to a subsequence $\{(\Phi_{tr}^{N'}, \Phi_e^{N'}, W^{N'}, I_e^{N'})\}$, which converges to a feasible quadruple $(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e)$ in the sense of Proposition 3.3. The lower semicontinuity of the objective follows as in [DACOROGNA 08], p. 96, Theorem 3.23., and p. 97, Remark 3.25.(ii). Consequently, denoting the minimal value of (P) by m , we get

$$\begin{aligned} m &= \lim_{N' \rightarrow \infty} F(\Phi_{tr}^{N'}, \Phi_e^{N'}, W^{N'}, I_e^{N'}) \\ &\geq \liminf_{N' \rightarrow \infty} F(\Phi_{tr}^{N'}, \Phi_e^{N'}, W^{N'}, I_e^{N'}) \geq F(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e) \geq m, \end{aligned} \tag{3.19}$$

and the quadruple $(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e)$ is a global minimizer of (P). ■

4. The adjoint equations.

4.1. Derivation of the adjoint system.

Throughout the following sections, we will further assume that the integrand $r(x, t, \varphi, \eta, w)$ within the objective (3.11) is continuously differentiable with respect to the variables φ , η and w . For the optimal control problem (P), let us introduce now the formal Lagrange function

$$\begin{aligned} \mathcal{L}(\Phi_{tr}, \Phi_e, W, I_e, P_1, P_2, P_3, P_4, P_5) &= F(\Phi_{tr}, \Phi_e, W, I_e) + \langle P_1, E_1(\Phi_{tr}, \Phi_e, W) \rangle \\ &+ \langle P_2, E_2(\Phi_{tr}, \Phi_e, I_e) \rangle + \langle P_3, E_3(\Phi_{tr}, W) \rangle + \langle P_4, E_4(\Phi_{tr}) \rangle + \langle P_5, E_5(W) \rangle \end{aligned} \quad (4.1)$$

with multipliers

$$P_1 \in L^4[(0, T), W^{1,2}(\Omega)]; \quad (4.2)$$

$$P_2 \in L^2[(0, T), W^{1,2}(\Omega)] \cap \left\{ Z \mid \int_{\Omega} Z(x, t) dx = 0 \quad (\forall) t \in (0, T) \right\}; \quad (4.3)$$

$$P_3 \in L^2[(0, T), L^2(\Omega)]; \quad P_4, P_5 \in (L^2(\Omega))^*. \quad (4.4)$$

Differentiating \mathcal{L} at the point $(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e)$ in a formal way with respect to the variables Φ_{tr} , Φ_e and W , we find the adjoint equations

$$\begin{aligned} D_{\Phi_{tr}} F(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e) + \langle P_1, D_{\Phi_{tr}} E_1(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \rangle \\ + \langle P_2, D_{\Phi_{tr}} E_2(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{I}_e) \rangle + \langle P_3, D_{\Phi_{tr}} E_3(\hat{\Phi}_{tr}, \hat{W}) \rangle + \langle P_4, D_{\Phi_{tr}} E_4(\hat{\Phi}_{tr}) \rangle = 0; \end{aligned} \quad (4.5)$$

$$D_{\Phi_e} F(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e) + \langle P_1, D_{\Phi_e} E_1(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \rangle + \langle P_2, D_{\Phi_e} E_2(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{I}_e) \rangle = 0; \quad (4.6)$$

$$\begin{aligned} D_W F(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e) + \langle P_1, D_W E_1(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \rangle + \langle P_3, D_W E_3(\hat{\Phi}_{tr}, \hat{W}) \rangle + \langle P_5, D_W E_5(\hat{W}) \rangle = 0. \end{aligned} \quad (4.7)$$

After choosing $P_4 = -P_1(\cdot, 0)$ and $P_5 = -P_3(\cdot, 0)$ (this choice is possible by Theorem 4.2. below), the adjoint system takes the following form:

$$\begin{aligned} \int_0^T \int_{\Omega} \left(-\frac{\partial P_1}{\partial t} + \frac{\partial I_{ion}}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{W}) P_1 + \frac{\partial G}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{W}) P_3 \right) \psi dx dt + \int_0^T \int_{\Omega} \nabla \psi^T M_i (\nabla P_1 + \nabla P_2) dx dt \\ = - \int_0^T \int_{\Omega} \left(\frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right) \psi dx dt \quad \forall \psi \in L^2[(0, T), W^{1,2}(\Omega)], \quad P_1(x, T) \equiv 0; \end{aligned} \quad (4.8)$$

$$\int_0^T \int_{\Omega} \nabla \psi^T M_i \nabla P_1 dx dt + \int_0^T \int_{\Omega} \nabla \psi^T (M_i + M_e) \nabla P_2 dx dt = - \int_0^T \int_{\Omega} \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \psi dx dt \quad (4.9)$$

$$\forall \psi \in L^2[(0, T), W^{1,2}(\Omega)] \quad \text{with} \quad \int_{\Omega} \psi(x, t) dx = 0 \quad (\forall) t \in (0, T), \quad \int_{\Omega} P_2(x, t) dx = 0 \quad (\forall) t \in (0, T);$$

$$\begin{aligned} \int_0^T \int_{\Omega} \left(-\frac{\partial P_3}{\partial t} + \frac{\partial I_{ion}}{\partial w}(\hat{\Phi}_{tr}, \hat{W}) P_1 + \frac{\partial G}{\partial w}(\hat{\Phi}_{tr}, \hat{W}) P_3 \right) \psi dx dt = - \int_0^T \int_{\Omega} \left(\frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right) \psi dx dt \\ \forall \psi \in L^2[(0, T), L^2(\Omega)], \quad P_3(x, T) \equiv 0. \end{aligned} \quad (4.10)$$

4.2. The reduced form of the adjoint system.

First, we apply to the system (4.8) – (4.10) the transformation $s = T - t$, thus defining $\widetilde{P}_i(x, s) = P_i(x, T - s)$, $1 \leq i \leq 3$, $\widetilde{\Phi}_{tr}(x, s) = \hat{\Phi}_{tr}(x, T - s)$, $\widetilde{\Phi}_e(x, s) = \hat{\Phi}_e(x, T - s)$, $\widetilde{W}(x, s) = \hat{W}(x, T - s)$ and $\widetilde{I}_e(x, s) = \hat{I}_e(x, T - s)$ etc. By abuse of notation, we suppress all tildes, thus simply replacing t by s and $-\partial P_1/\partial t$, $-\partial P_3/\partial t$ by $\partial P_1/\partial s$ and $\partial P_3/\partial s$, respectively. Then the adjoint system, in analogy to the primal bidomain equations, can be rewritten in terms of the bidomain bilinear form as a reduced system:

$$\frac{d}{ds} \langle P_1(s), \psi \rangle + A(P_1(s), \psi) + \int_{\Omega} \left(\frac{\partial I_{ion}}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{W}) P_1 + \frac{\partial G}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{W}) P_3 \right) \psi dx = \langle \widetilde{S}(s), \psi \rangle \quad (4.11)$$

$\forall \psi \in W^{1,2}(\Omega);$

$$\frac{d}{ds} \langle P_3(s), \psi \rangle + \int_{\Omega} \left(\frac{\partial I_{ion}}{\partial w}(\hat{\Phi}_{tr}, \hat{W}) P_1 + \frac{\partial G}{\partial w}(\hat{\Phi}_{tr}, \hat{W}) P_3 \right) \psi dx = - \left\langle \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}), \psi \right\rangle \quad (4.12)$$

$\forall \psi \in L^2(\Omega);$

$$P_1(x, 0) = 0 \quad (\forall) x \in \Omega; \quad P_3(x, 0) = 0 \quad (\forall) x \in \Omega \quad (4.13)$$

on $[0, T]$ in distributional sense, cf. [KUNISCH/WAGNER 11], p. 5 f., Theorem 2.4. Here the bidomain bilinear form $A: W^{1,2}(\Omega) \times W^{1,2}(\Omega) \rightarrow \mathbb{R}$ is defined as ibid., p. 5, (2.22), through

$$A(\psi_1, \psi_2) = \int_{\Omega} \nabla \psi_1^T M_i \nabla \psi_2 dx + \int_{\Omega} \nabla \widetilde{\psi}_e^T M_i \nabla \psi_2 dx \quad (4.14)$$

where $\widetilde{\psi}_e \in W^{1,2}(\Omega)$ is the uniquely determined solution of the variational equation

$$\int_{\Omega} \nabla \widetilde{\psi}_e^T (M_i + M_e) \nabla \psi dx = - \int_{\Omega} \nabla \psi_1^T M_i \nabla \psi dx \quad \forall \psi \in W^{1,2}(\Omega) \quad \text{with} \quad \int_{\Omega} \psi dx = 0 \quad (4.15)$$

satisfying $\int_{\Omega} \widetilde{\psi}_e dx = 0,$

and the linear functionals $\widetilde{S}(s) \in (W^{1,2}(\Omega))^*$ are defined through

$$\langle \widetilde{S}(s), \psi \rangle = - \left\langle \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}), \psi \right\rangle - \int_{\Omega} \nabla \bar{\psi}_e^T M_i \nabla \psi dx \quad (4.16)$$

where $\bar{\psi}_e \in W^{1,2}(\Omega)$ is the uniquely determined solution of the variational equation

$$\int_{\Omega} \nabla \bar{\psi}_e^T (M_i + M_e) \nabla \psi dx = \left\langle \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}), \psi \right\rangle \quad \forall \psi \in W^{1,2}(\Omega) \quad \text{with} \quad \int_{\Omega} \psi dx = 0 \quad (4.17)$$

satisfying $\int_{\Omega} \bar{\psi}_e dx = 0.$

The component P_2 of the solution of (4.8) – (4.10) is uniquely determined as the sum $P_2 = \widetilde{\psi}_e + \bar{\psi}_e$. Note that this reformulation is even possible without imposing the additional compatibility condition

$$\int_{\Omega} \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}(x, s), \hat{\Phi}_e(x, s), \hat{W}(x, s)) dx = 0 \quad (\forall) s \in (0, T). \quad (4.18)$$

4.3. Existence and regularity of weak solutions.

Theorem 4.1. (A-priori estimates for weak solutions of the adjoint system) *The optimal control problem (3.11) – (3.17) is studied under the assumptions from Subsection 3.1. Within the problem, we specify*

the Rogers-McCulloch model. Assume further that the integrand $r(x, t, \varphi, \eta, w)$ is continuously differentiable with respect to φ, η and w .

1) If $(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e)$ is a feasible solution of (P) with

$$\frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}), \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}), \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \in L^2(\Omega_T) \quad (4.19)$$

then every weak solution $(P_1, P_2, P_3) \in L^2[(0, T), W^{1,2}(\Omega)] \times L^2[(0, T), W^{1,2}(\Omega)] \times L^2(\Omega_T)$ of the adjoint system (4.8) – (4.10) obeys the estimate

$$\begin{aligned} & \|P_1\|_{L^\infty[(0, T), L^2(\Omega)]}^2 + \|P_1\|_{L^2[(0, T), W^{1,2}(\Omega)]}^2 + \|P_2\|_{L^2[(0, T), W^{1,2}(\Omega)]}^2 + \|P_3\|_{L^\infty[(0, T), L^2(\Omega)]}^2 \\ & \leq C \left(\left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 + \left\| \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 \right) \end{aligned} \quad (4.20)$$

where the constant $C > 0$ does not depend on P_1, P_2, P_3 but on $(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e)$ and the data of (P).

2) Let $q = 10/9$. If $(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e)$ is a feasible solution of (P) with

$$\frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}), \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}), \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \in L^{2q}[(0, T), L^2(\Omega)] \quad (4.21)$$

then every weak solution $(P_1, P_2, P_3) \in L^2[(0, T), W^{1,2}(\Omega)] \times L^2[(0, T), W^{1,2}(\Omega)] \times L^2(\Omega_T)$ of the adjoint system (4.8) – (4.10) obeys (4.20) as well as the further estimate

$$\begin{aligned} & \|P_1\|_{C^0[0, T], L^2(\Omega)}^2 + \|\partial P_1 / \partial s\|_{L^q[(0, T), (W^{1,2}(\Omega))^*]}^q + \|P_3\|_{C^0[0, T], L^2(\Omega)}^2 \\ & + \|\partial P_3 / \partial s\|_{L^q[(0, T), L^2(\Omega)]}^q \leq \left(1 + \left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^{2q}[(0, T), L^2(\Omega)]}^{2q} \right. \\ & \left. + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^{2q}[(0, T), L^2(\Omega)]}^{2q} + \left\| \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 \right) \end{aligned} \quad (4.22)$$

where the constant $C > 0$ does not depend on P_1, P_2, P_3 but on $(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e)$ and the data of (P).

The a-priori estimates yield the following existence and uniqueness theorem for the adjoint system:

Theorem 4.2. (Existence and uniqueness of weak solutions for the adjoint system) *Under the assumptions of Theorem 4.1., 2), the adjoint system (4.8) – (4.10) admits a uniquely determined weak solution (P_1, P_2, P_3) with*

$$P_1 \in C^0[0, T], L^2(\Omega) \cap L^2[(0, T), W^{1,2}(\Omega)] \cap W^{1,q}[(0, T), (W^{1,2}(\Omega))^*]; \quad (4.23)$$

$$P_2 \in L^2[(0, T), W^{1,2}(\Omega)]; \quad \int_{\Omega} P_2(x, t) dx = 0 \quad (\forall) t \in (0, T); \quad (4.24)$$

$$P_3 \in C^0[0, T], L^2(\Omega) \cap W^{1,q}[(0, T), L^2(\Omega)]. \quad (4.25)$$

Note that, even under the assumptions of Theorems 4.1., 2) and 4.2., the regularity of $P_1 \in L^4[(0, T), W^{1,2}(\Omega)]$ as required in (4.1) and (4.2) cannot be guaranteed.

4.4. Proofs.

Proof of Theorem 4.1. Throughout the proof, C denotes a generical positive constant, which may appropriately change from line to line. Further, we will specify in (3.12) – (3.14) the Rogers-McCulloch model. The necessary alterations in the case of the other models will be discussed at the end of the subsection.

• **Step 1.** An estimate for the right-hand side of (4.11). We start with

Lemma 4.4. Under the assumptions of Theorem 4.1., for arbitrary $\varepsilon'_0 > 0$ the following estimate holds:

$$|\langle \tilde{S}(s), \psi \rangle| \leq \frac{C}{\varepsilon'_0} \left(\left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2 + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2 \right) + C \varepsilon'_0 \|\psi\|_{W^{1,2}(\Omega)}^2. \quad (4.26)$$

The constant $C > 0$ does not depend on ε'_0 and ψ .

Proof. Inserting $\bar{\psi}_e \in W^{1,2}(\Omega)$ as a feasible test function into (4.17), we get from the uniform ellipticity of M_i and M_e and the Poincaré inequality:

$$\begin{aligned} C \|\bar{\psi}_e\|_{W^{1,2}(\Omega)}^2 &\leq \int_{\Omega} \nabla \bar{\psi}_e^T (M_i + M_e) \nabla \bar{\psi}_e^T dx \leq |\langle \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}), \bar{\psi}_e^T \rangle| \\ &\leq \frac{1}{2\delta_1} \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{(W^{1,2}(\Omega))^*}^2 + \frac{\delta_1}{2} \|\bar{\psi}_e\|_{W^{1,2}(\Omega)}^2 \quad (\forall) s \in (0, T), \end{aligned} \quad (4.27)$$

for arbitrary $\delta_1 > 0$. Inserting $\delta_1 = C$, we arrive at

$$\frac{C}{2} \|\bar{\psi}_e\|_{W^{1,2}(\Omega)}^2 \leq \frac{1}{2C} \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{(W^{1,2}(\Omega))^*}^2 \leq \frac{1}{2C} \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2. \quad (4.28)$$

From (4.16), we obtain

$$|\langle \tilde{S}(s), \psi \rangle| \leq |\langle \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}), \psi \rangle| + |\langle \nabla \bar{\psi}_e^T M_i, \nabla \psi \rangle| \quad (4.29)$$

$$\leq \frac{1}{2\delta_2} \left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{(W^{1,2}(\Omega))^*}^2 + \frac{\delta_2}{2} \|\psi\|_{W^{1,2}(\Omega)}^2 + \frac{1}{2\delta_3} \|\bar{\psi}_e\|_{L^2(\Omega)}^2 + \frac{\delta_3}{2} \|M_i\|^2 \cdot \|\psi\|_{L^2(\Omega)}^2 \quad (4.30)$$

$$\leq \frac{1}{2\delta_2} \left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2 + \frac{1}{2\delta_3} \|\bar{\psi}_e\|_{W^{1,2}(\Omega)}^2 + \left(\frac{\delta_2}{2} + \frac{(\mu_2)^2 \delta_3}{2} \right) \|\psi\|_{W^{1,2}(\Omega)}^2 \quad (4.31)$$

$$\leq \frac{C}{\delta_2} \left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2 + \frac{C}{\delta_3} \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2 + C \left(\frac{\delta_2}{2} + \frac{(\mu_2)^2 \delta_3}{2} \right) \|\psi\|_{W^{1,2}(\Omega)}^2 \quad (4.32)$$

by (2.5) and (4.28). Taking $\delta_3 = \delta_2/(\mu_2)^2$, we get (4.26). ■

• **Step 2.** The estimates for $\|P_1\|_{L^\infty[0, T], L^2(\Omega)}$ and $\|P_3\|_{L^\infty[0, T], L^2(\Omega)}$. Specifying the derivatives of I_{ion} and G according to the Rogers-McCulloch model, we have

$$\frac{\partial I_{ion}}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{W}) = 3b(\hat{\Phi}_{tr})^2 - 2(a+1)b\hat{\Phi}_{tr} + ab + \hat{W}; \quad \frac{\partial G}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{W}) \equiv -\varepsilon\kappa; \quad (4.33)$$

$$\frac{\partial I_{ion}}{\partial w}(\hat{\Phi}_{tr}, \hat{W}) = \hat{\Phi}_{tr}; \quad \frac{\partial G}{\partial w}(\hat{\Phi}_{tr}, \hat{W}) \equiv \varepsilon. \quad (4.34)$$

Inserting $P_1(s)$ as a feasible test function into (4.11), we get for arbitrary $\varepsilon'_0, \varepsilon_1(s) > 0$ with [KUNISCH/WAGNER 11], p. 6, Theorem 2.4., 2), and Lemma 4.3. above¹¹⁾

$$\frac{1}{2} \frac{d}{ds} \|P_1(s)\|_{L^2(\Omega)}^2 + \left(A(P_1, P_1) + \beta \|P_1\|_{L^2(\Omega)}^2 \right) + \int_{\Omega} \frac{\partial I_{ion}}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{W}) P_1(s)^2 dx \quad (4.35)$$

$$\leq \int_{\Omega} \left| \frac{\partial G}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{W}) \right| |P_1 P_3| + |\langle \tilde{S}(s), P_1 \rangle| + \beta \|P_1\|_{L^2(\Omega)}^2 \implies$$

$$\frac{1}{2} \frac{d}{ds} \|P_1(s)\|_{L^2(\Omega)}^2 + \beta \|P_1\|_{W^{1,2}(\Omega)}^2 + \int_{\Omega} (ab + 3b(\hat{\Phi}_{tr})^2) P_1(s)^2 dx \quad (4.36)$$

$$\leq C \int_{\Omega} (|\hat{\Phi}_{tr}| + |\hat{W}|) |P_1|^2 dx + \varepsilon\kappa \int_{\Omega} |P_1 P_3| dx$$

$$+ \frac{C}{\varepsilon'_0} \left(\left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2 + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2 \right) + C \varepsilon'_0 \|P_1\|_{W^{1,2}(\Omega)}^2 + \beta \|P_1\|_{L^2(\Omega)}^2 \implies$$

¹¹⁾ Note that $\varepsilon > 0$ is fixed from the Rogers-McCulloch model.

$$\frac{1}{2} \frac{d}{ds} \|P_1(s)\|_{L^2(\Omega)}^2 + \beta \|P_1\|_{W^{1,2}(\Omega)}^2 \quad (4.37)$$

$$\begin{aligned} &\leq C \varepsilon_1(s) \int_{\Omega} \left(|\hat{\Phi}_{tr}|^2 + |\hat{W}|^2 \right) |P_1|^2 dx + \frac{C}{\varepsilon_1(s)} \|P_1\|_{L^2(\Omega)}^2 + C \left(\|P_1\|_{L^2(\Omega)}^2 + \|P_3\|_{L^2(\Omega)}^2 \right) \\ &+ \frac{C}{\varepsilon_0} \left(\left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2 + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2 \right) + C \varepsilon_0' \|P_1\|_{W^{1,2}(\Omega)}^2 + \beta \|P_1\|_{L^2(\Omega)}^2 \\ &\leq C \varepsilon_1(s) \left(\|\hat{\Phi}_{tr}\|_{L^4(\Omega)}^2 + \|\hat{W}\|_{L^4(\Omega)}^2 \right) \cdot \|P_1\|_{W^{1,2}(\Omega)}^2 + C \left(1 + \frac{1}{\varepsilon_1(s)} \right) \|P_1\|_{L^2(\Omega)}^2 \\ &+ \frac{C}{\varepsilon_0} \left(\left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2 + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2 \right) + C \varepsilon_0' \|P_1\|_{W^{1,2}(\Omega)}^2 + C \|P_3\|_{L^2(\Omega)}^2. \end{aligned} \quad (4.38)$$

We choose $\varepsilon_1(s) = \varepsilon_1' / (1 + \|\hat{\Phi}_{tr}(s)\|_{L^4(\Omega)}^2 + \|\hat{W}(s)\|_{L^4(\Omega)}^2)$ with $\varepsilon_1' > 0$ and continue (4.38) with

$$\begin{aligned} \dots &\leq C \varepsilon_1' \cdot \|P_1\|_{W^{1,2}(\Omega)}^2 + C \left(1 + \frac{1}{\varepsilon_1'} (1 + \|\hat{\Phi}_{tr}(s)\|_{L^4(\Omega)}^2 + \|\hat{W}(s)\|_{L^4(\Omega)}^2) \right) \cdot \|P_1\|_{L^2(\Omega)}^2 \\ &+ \frac{C}{\varepsilon_0'} \left(\left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2 + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2 \right) + C \varepsilon_0' \|P_1\|_{W^{1,2}(\Omega)}^2 + C \|P_3\|_{L^2(\Omega)}^2. \end{aligned} \quad (4.39)$$

Further, inserting $P_3(s)$ as a feasible test function into (4.12), we find with $\varepsilon_2(s) > 0$

$$\begin{aligned} \frac{1}{2} \frac{d}{ds} \|P_3(s)\|_{L^2(\Omega)}^2 &\leq \int_{\Omega} \left| \frac{\partial I_{ion}}{\partial w}(\hat{\Phi}_{tr}, \hat{W}) \right| |P_1 P_3| dx + \int_{\Omega} \left| \frac{\partial G}{\partial w}(\hat{\Phi}_{tr}, \hat{W}) \right| |P_3|^2 dx + \int_{\Omega} \left| \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right| |P_3| dx \end{aligned} \quad (4.40)$$

$$\leq \int_{\Omega} |\hat{\Phi}_{tr} P_1 P_3| dx + \varepsilon \|P_3\|_{L^2(\Omega)}^2 + C \left\| \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2 + C \|P_3\|_{L^2(\Omega)}^2 \quad (4.41)$$

$$\leq C \varepsilon_2(s) \int_{\Omega} |\hat{\Phi}_{tr} P_1|^2 dx + C \left(1 + \frac{1}{\varepsilon_2(s)} \right) \|P_3\|_{L^2(\Omega)}^2 + C \left\| \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2 \quad (4.42)$$

$$\leq C \varepsilon_2(s) \|\hat{\Phi}_{tr}\|_{L^4(\Omega)}^2 \cdot \|P_1\|_{W^{1,2}(\Omega)}^2 + C \left(1 + \frac{1}{\varepsilon_2(s)} \right) \|P_3\|_{L^2(\Omega)}^2 + C \left\| \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2. \quad (4.43)$$

Choosing now $\varepsilon_2(s) = \varepsilon_2' / (1 + \|\hat{\Phi}_{tr}(s)\|_{L^4(\Omega)}^2)$ with $\varepsilon_2' > 0$, (4.43) may be continued as

$$\begin{aligned} \dots &\leq C \varepsilon_2' \|P_1\|_{W^{1,2}(\Omega)}^2 + C \left(1 + \frac{1}{\varepsilon_2'} (1 + \|\hat{\Phi}_{tr}(s)\|_{L^4(\Omega)}^2) \right) \|P_3\|_{L^2(\Omega)}^2 \\ &+ C \left\| \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2. \end{aligned} \quad (4.44)$$

Combining (4.39) and (4.44), we obtain

$$\begin{aligned} \frac{d}{ds} \left(\|P_1(s)\|_{L^2(\Omega)}^2 + \|P_3(s)\|_{L^2(\Omega)}^2 \right) + 2\beta \|P_1\|_{W^{1,2}(\Omega)}^2 & \\ \leq C (\varepsilon_0' + \varepsilon_1' + \varepsilon_2') \|P_1\|_{W^{1,2}(\Omega)}^2 + C \left(1 + \frac{1}{\varepsilon_1'} (1 + \|\hat{\Phi}_{tr}(s)\|_{L^4(\Omega)}^2 + \|\hat{W}(s)\|_{L^4(\Omega)}^2) \right) \|P_1\|_{L^2(\Omega)}^2 & \\ + C \left(1 + \frac{1}{\varepsilon_2'} (1 + \|\hat{\Phi}_{tr}(s)\|_{L^4(\Omega)}^2) \right) \|P_3\|_{L^2(\Omega)}^2 & \\ + \frac{C}{\varepsilon_0'} \left(\left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2 + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2 \right) + C \left\| \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2. & \end{aligned} \quad (4.45)$$

Now we fix the parameters $\varepsilon_0', \varepsilon_1', \varepsilon_2' > 0$ in such a way that the terms with $\|P_1\|_{W^{1,2}(\Omega)}$ on both sides of (4.45) will be annihilated, thus arriving at

$$\frac{d}{ds} \left(\|P_1(s)\|_{L^2(\Omega)}^2 + \|P_3(s)\|_{L^2(\Omega)}^2 \right) \leq A(s) \left(\|P_1(s)\|_{L^2(\Omega)}^2 + \|P_3(s)\|_{L^2(\Omega)}^2 \right) + B(s) \quad \text{where} \quad (4.46)$$

$$A(s) = C \left(1 + \|\hat{\Phi}_{tr}(s)\|_{L^4(\Omega)}^2 + \|\hat{W}(s)\|_{L^4(\Omega)}^2 \right); \quad (4.47)$$

$$B(s) = C \left(\left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2 + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2 + \left\| \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2 \right). \quad (4.48)$$

Then Gronwall's inequality yields for all $s \in [0, T]$:

$$\|P_1(s)\|_{L^2(\Omega)}^2 + \|P_3(s)\|_{L^2(\Omega)}^2 \leq e^{\int_0^s A(\sigma) d\sigma} \left(\|P_1(0)\|_{L^2(\Omega)}^2 + \|P_3(0)\|_{L^2(\Omega)}^2 + \int_0^s B(\sigma) d\sigma \right) \quad (4.49)$$

$$\leq e^{CT} \left(\left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 + \left\| \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 \right) \quad (4.50)$$

since $\hat{\Phi}_{tr} \in L^4(\Omega_T)$ and $\hat{W} \in C^0[[0, T], L^4(\Omega)]$. Consequently, we get the estimate

$$\begin{aligned} \|P_1\|_{L^\infty[(0, T), L^2(\Omega)]}^2 + \|P_3\|_{L^\infty[(0, T), L^2(\Omega)]}^2 &\leq C \left(\left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 \right. \\ &\quad \left. + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 + \left\| \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 \right). \end{aligned} \quad (4.51)$$

• **Step 3.** *The estimate for $\|P_1\|_{L^2[(0, T), W^{1,2}(\Omega)]}^2$.* We return to (4.45). Then $\varepsilon'_0, \varepsilon'_1, \varepsilon'_2 > 0$ may be alternatively chosen in such a way that $C(\varepsilon'_0 + \varepsilon'_1 + \varepsilon'_2) = \beta$ and, consequently,

$$\begin{aligned} \frac{d}{ds} \left(\|P_1(s)\|_{L^2(\Omega)}^2 + \|P_3(s)\|_{L^2(\Omega)}^2 \right) + \beta \|P_1(s)\|_{W^{1,2}(\Omega)}^2 &\leq A(s) \left(\|P_1(s)\|_{L^2(\Omega)}^2 + \|P_3(s)\|_{L^2(\Omega)}^2 \right) + B(s) \end{aligned} \quad (4.52)$$

where $A(s)$ and $B(s)$ are calculated as above. Together with (4.51), we obtain

$$\begin{aligned} \frac{d}{ds} \left(\|P_1(s)\|_{L^2(\Omega)}^2 + \|P_3(s)\|_{L^2(\Omega)}^2 \right) + \beta \|P_1(s)\|_{W^{1,2}(\Omega)}^2 &\leq C A(s) \left(\left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 + \left\| \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 \right) + B(s). \end{aligned} \quad (4.53)$$

We integrate (4.53) over $[0, T]$ and get, inserting the initial values $P_1(0) = \mathbf{o}, P_3(0) = \mathbf{o}$

$$\|P_1(T)\|_{L^2(\Omega)}^2 + \|P_3(T)\|_{L^2(\Omega)}^2 + \beta \|P_1\|_{L^2[(0, T), W^{1,2}(\Omega)]}^2 \quad (4.54)$$

$$\leq C \left(\left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 + \left\| \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 \right) \implies$$

$$\|P_1\|_{L^2[(0, T), W^{1,2}(\Omega)]}^2 \quad (4.55)$$

$$\leq C \left(\left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 + \left\| \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 \right).$$

• **Step 4.** *The estimate for $\|\partial P_1/\partial s\|_{L^q[(0, T), (W^{1,2}(\Omega))^*]}^2$ with $q = 10/9 < 2$.* Exploiting the definition of the dual norm, we start with

$$\|\partial P_1/\partial s\|_{L^q[(0, T), (W^{1,2}(\Omega))^*]}^q = \int_0^T \sup_{\|\psi\|_{W^{1,2}(\Omega)}=1} |\langle \partial P_1(s)/\partial s, \psi \rangle|^q ds \quad (4.56)$$

$$= \int_0^T \sup_{\dots} \left| -A(P_1, \psi) - \int_{\Omega} \left(\frac{\partial I_{ion}}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{W}) P_1 + \frac{\partial G}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{W}) P_3 \right) \psi dx + \langle \tilde{S}(s), \psi \rangle \right|^q ds \quad (4.57)$$

$$\begin{aligned} \leq C \int_0^T \left(\sup_{\dots} |A(P_1, \psi)|^q + \sup_{\dots} \left(\int_{\Omega} \left(|\hat{\Phi}_{tr}|^2 + |\hat{\Phi}_{tr}| + |\hat{W}| + 1 \right) |P_1| |\psi| dx \right)^q \right. \\ \left. + \sup_{\dots} \varepsilon^q \kappa^q \left(\int_{\Omega} |P_3| |\psi| dx \right)^q + \sup_{\dots} |\langle \tilde{S}(s), \psi \rangle|^q \right) ds. \end{aligned} \quad (4.58)$$

The four terms on the right-hand side of (4.58) will be estimated separately. For the first term, we get with [KUNISCH/WAGNER 11], p. 6, Theorem 2.4., 2), and (4.55)

$$\int_0^T \sup_{\dots} |A(P_1, \psi)|^q ds \leq \int_0^T \sup_{\dots} \gamma^q \|P_1(s)\|_{W^{1,2}(\Omega)}^q \|\psi\|_{W^{1,2}(\Omega)}^q ds \quad (4.59)$$

$$\leq \gamma^q \|P_1\|_{L^q[(0,T), W^{1,2}(\Omega)]}^q \leq C \|P_1\|_{L^2[(0,T), W^{1,2}(\Omega)]}^q \quad (4.60)$$

$$\leq C \left(\left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 + \left\| \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 \right)^{q/2}. \quad (4.61)$$

For the second term, we obtain

$$\int_0^T \sup_{\dots} \left(\int_{\Omega} (|\hat{\Phi}_{tr}|^2 + |\hat{\Phi}_{tr}| + |\hat{W}| + 1) |P_1| |\psi| dx \right)^q ds \quad (4.62)$$

$$\leq C \left(\sup_{\dots} \int_0^T \int_{\Omega} |\hat{\Phi}_{tr}|^{2q} |P_1|^q |\psi|^q dx ds + \sup_{\dots} \int_0^T \int_{\Omega} |\hat{\Phi}_{tr}|^q |P_1|^q |\psi|^q dx ds \right. \\ \left. + \sup_{\dots} \int_0^T \int_{\Omega} |\hat{W}|^q |P_1|^q |\psi|^q dx ds + \sup_{\dots} \int_0^T \int_{\Omega} |P_1|^q |\psi|^q dx ds \right) = J_1 + J_2 + J_3 + J_4. \quad (4.63)$$

We start with the estimation of J_1 , thus getting

$$J_1 = \sup_{\dots} \int_0^T \int_{\Omega} |\hat{\Phi}_{tr}|^{20/9} |P_1|^{10/9} |\psi|^{10/9} dx ds \quad (4.64)$$

$$= \sup_{\dots} \int_0^T \left(\int_{\Omega} |\hat{\Phi}_{tr}|^{40/11} dx \right)^{33/54} \left(\int_{\Omega} |P_1|^{60/11} dx \right)^{11/54} \left(\int_{\Omega} |\psi|^6 dx \right)^{10/54} ds \quad (4.65)$$

$$= \sup_{\dots} \int_0^T \|\hat{\Phi}_{tr}\|_{L^{40/11}(\Omega)}^{20/9} \cdot \|P_1\|_{L^{60/11}(\Omega)}^{10/9} \cdot \|\psi\|_{L^6(\Omega)}^{10/9} ds \quad (4.66)$$

$$\leq \sup_{\dots} C \int_0^T \|\hat{\Phi}_{tr}\|_{L^{40/11}(\Omega)}^{20/9} \cdot \|P_1\|_{W^{1,2}(\Omega)}^{10/9} \cdot \|\psi\|_{W^{1,2}(\Omega)}^{10/9} ds = C \int_0^T \|\hat{\Phi}_{tr}\|_{L^{40/11}(\Omega)}^{20/9} \cdot \|P_1\|_{W^{1,2}(\Omega)}^{10/9} ds \quad (4.67)$$

$$\leq C \left(\int_0^T \|\hat{\Phi}_{tr}(s)\|_{L^{40/11}(\Omega)}^5 ds \right)^{8/18} \left(\int_0^T \|P_1(s)\|_{W^{1,2}(\Omega)}^2 ds \right)^{10/18} \quad (4.68)$$

$$\leq C \cdot \|\hat{\Phi}_{tr}\|_{L^5[(0,T), L^{40/11}(\Omega)]}^{20/9} \cdot \|P_1\|_{L^2[(0,T), W^{1,2}(\Omega)]}^{10/9}. \quad (4.69)$$

Since $\hat{\Phi}_{tr} \in L^{p'}[(0,T), L^{p''}(\Omega)]$ for all $1 < p' < \infty$, $4 \leq p'' < 6$, we get a bound analogous to (4.61). Continuing with J_2 , we find in completely analogous manner

$$J_2 = \sup_{\dots} \int_0^T \int_{\Omega} |\hat{\Phi}_{tr}|^{10/9} |P_1|^{10/9} |\psi|^{10/9} dx ds \quad (4.70)$$

$$\leq \sup_{\dots} C \int_0^T \|\hat{\Phi}_{tr}\|_{L^{20/11}(\Omega)}^{10/9} \cdot \|P_1\|_{W^{1,2}(\Omega)}^{10/9} \cdot \|\psi\|_{W^{1,2}(\Omega)}^{10/9} ds = C \int_0^T \|\hat{\Phi}_{tr}\|_{L^{20/11}(\Omega)}^{10/9} \cdot \|P_1\|_{W^{1,2}(\Omega)}^{10/9} ds \quad (4.71)$$

$$\leq C \left(\int_0^T \|\hat{\Phi}_{tr}(s)\|_{L^{20/11}(\Omega)}^{10/4} ds \right)^{8/18} \left(\int_0^T \|P_1(s)\|_{W^{1,2}(\Omega)}^2 ds \right)^{10/18} \quad (4.72)$$

$$\leq C \cdot \|\hat{\Phi}_{tr}\|_{L^{10/4}[(0,T), L^{20/11}(\Omega)]}^{10/9} \cdot \|P_1\|_{L^2[(0,T), W^{1,2}(\Omega)]}^{10/9}. \quad (4.73)$$

Noticing that, in the case of the Rogers-McCulloch model, \hat{W} belongs even to $C^0[[0,T], L^4(\Omega)]$, we may

estimate J_3 in the same way:

$$J_3 = \sup_{\dots} \int_0^T \int_{\Omega} |\hat{W}|^{10/9} |P_1|^{10/9} |\psi|^{10/9} dx ds \quad (4.74)$$

$$\leq \sup_{\dots} C \int_0^T \|\hat{W}\|_{L^{20/11}(\Omega)}^{10/9} \cdot \|P_1\|_{W^{1,2}(\Omega)}^{10/9} \cdot \|\psi\|_{W^{1,2}(\Omega)}^{10/9} ds = C \int_0^T \|\hat{W}\|_{L^{20/11}(\Omega)}^{10/9} \cdot \|P_1\|_{W^{1,2}(\Omega)}^{10/9} ds \quad (4.75)$$

$$\leq C \left(\int_0^T \|\hat{W}(s)\|_{L^{20/11}(\Omega)}^{10/4} ds \right)^{8/18} \left(\int_0^T \|P_1(s)\|_{W^{1,2}(\Omega)}^2 ds \right)^{10/18} \quad (4.76)$$

$$\leq C \cdot \|\hat{W}\|_{L^{10/4}[(0,T), L^{20/11}(\Omega)]}^{10/9} \cdot \|P_1\|_{L^2[(0,T), W^{1,2}(\Omega)]}^{10/9} \quad (4.77)$$

$$\leq C \cdot \|\hat{W}\|_{L^\infty[(0,T), L^4(\Omega)]}^{10/9} \cdot \|P_1\|_{L^2[(0,T), W^{1,2}(\Omega)]}^{10/9} \quad (4.78)$$

Defining the function $S(x, s) \equiv 1$, the estimation of J_4 yields

$$J_4 = \sup_{\dots} \int_0^T \int_{\Omega} |S|^{10/9} |P_1|^{10/9} |\psi|^{10/9} dx ds \quad (4.79)$$

$$\leq C \cdot \|S\|_{L^{10/4}[(0,T), L^{20/11}(\Omega)]}^{10/9} \cdot \|P_1\|_{L^2[(0,T), W^{1,2}(\Omega)]}^{10/9}.$$

Summing up, we get from (4.63), (4.69), (4.73), (4.78) and (4.79):

$$\int_0^T \sup_{\dots} \left(\int_{\Omega} (|\hat{\Phi}_{tr}|^2 + |\hat{\Phi}_{tr}| + |\hat{W}| + 1) |P_1| |\psi| dx \right)^q ds \quad (4.80)$$

$$\leq C \left(\left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 + \left\| \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 \right)^{q/2}.$$

The third term at the right-hand side of (4.58) will be estimated through

$$\int_0^T \sup_{\dots} \varepsilon^q \kappa^q \left(\int_{\Omega} |P_3| |\psi| dx \right)^q ds \leq \sup_{\dots} C \int_0^T \int_{\Omega} |P_3|^{10/9} |\psi|^{10/9} dx ds \quad (4.81)$$

$$\leq \sup_{\dots} C \int_0^T \left(\int_{\Omega} |P_3|^2 dx \right)^{10/18} \left(\int_{\Omega} |\psi|^{10/4} dx \right)^{8/18} ds = \sup_{\dots} C \int_0^T \|P_3\|_{L^2(\Omega)}^{10/9} \cdot \|\psi\|_{L^{10/4}(\Omega)}^{10/9} ds \quad (4.82)$$

$$\leq \sup_{\dots} C \int_0^T \|P_3\|_{L^2(\Omega)}^{10/9} \cdot \|\psi\|_{W^{1,2}(\Omega)}^{10/9} = C \|P_3\|_{L^q[(0,T), L^2(\Omega)]}^q \leq C \|P_3\|_{L^\infty[(0,T), L^2(\Omega)]}^q \quad (4.83)$$

$$\leq C \left(\left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 + \left\| \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 \right)^{q/2}. \quad (4.84)$$

Finally, Lemma 4.3. implies for the fourth term at the right-hand side of (4.58):

$$\int_0^T \sup_{\dots} |\langle \tilde{S}(s), \psi \rangle|^q ds \leq C \left(1 + \int_0^T \left(\left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2 + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2 \right)^q ds \right) \quad (4.85)$$

$$\leq C \left(1 + \int_0^T \left(\left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^{2q} + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^{2q} \right) ds \right) \quad (4.86)$$

$$\leq C \left(1 + \left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^{20/9}[(0,T), L^2(\Omega)]}^{20/9} + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^{20/9}[(0,T), L^2(\Omega)]}^{20/9} \right). \quad (4.87)$$

Together with (4.61), (4.80), (4.84) and (4.87), (4.58) yields the claimed estimate

$$\|\partial P_1 / \partial s\|_{L^q[(0,T), (W^{1,2}(\Omega))^*]}^q \quad (4.88)$$

$$\leq C \left(\left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 + \left\| \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 \right)^{q/2}$$

$$+ C \left(1 + \left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^{20/9}[(0,T), L^2(\Omega)]}^{20/9} + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^{20/9}[(0,T), L^2(\Omega)]}^{20/9} \right).$$

• **Step 5.** The estimate for $\|\partial P_3/\partial s\|_{L^q[(0,T), (L^2(\Omega))^*]}^q$. We start again by using the dual norm

$$\|\partial P_3/\partial s\|_{L^q[(0,T), (L^2(\Omega))^*]}^q = \int_0^T \sup_{\|\psi\|_{L^2(\Omega)}=1} |\langle \partial P_3(s)/\partial s, \psi \rangle|^q ds \quad (4.89)$$

$$= \int_0^T \sup_{\dots} \left| \int_{\Omega} \left(-\hat{\Phi}_{tr} P_1 - \varepsilon P_3 - \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right) \psi dx \right|^q ds \quad (4.90)$$

$$\begin{aligned} &\leq C \int_0^T \sup_{\dots} \left(\left(\int_{\Omega} |\hat{\Phi}_{tr}| |P_1| |\psi| dx \right)^q + \left(\varepsilon \int_{\Omega} |P_3| |\psi| dx \right)^q + \left(\int_{\Omega} \left| \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right| |\psi| dx \right)^q \right) ds \\ &\leq C \left(\sup_{\dots} \int_0^T \int_{\Omega} |\hat{\Phi}_{tr}|^q |P_1|^q |\psi|^q dx ds + \sup_{\dots} \int_0^T \left(\int_{\Omega} |P_3| |\psi| dx \right)^q ds \right. \\ &\quad \left. + \sup_{\dots} \int_0^T \left(\int_{\Omega} \left| \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right| |\psi| dx \right)^q ds \right) = J_5 + J_6 + J_7. \end{aligned} \quad (4.91)$$

The three terms on the right-hand side of (4.91) will be estimated separately. For the first term, we get

$$J_5 = \sup_{\dots} C \int_0^T \int_{\Omega} |\hat{\Phi}_{tr}|^{10/9} |P_1|^{10/9} |\psi|^{10/9} dx ds \quad (4.92)$$

$$\leq \sup_{\dots} C \left(\int_0^T \int_{\Omega} |\hat{\Phi}_{tr}|^5 dx ds \right)^{2/9} \left(\int_0^T \int_{\Omega} |P_1|^5 dx ds \right)^{2/9} \left(\int_0^T \int_{\Omega} |\psi|^2 dx ds \right)^{5/9} \quad (4.93)$$

$$= \sup_{\|\psi(s)\|_{L^2(\Omega)}=1} C \cdot \|\hat{\Phi}_{tr}\|_{L^5(\Omega_T)}^{10/9} \cdot \|P_1\|_{L^5(\Omega_T)}^{10/9} \cdot \left(\int_0^T \|\psi(s)\|_{L^2(\Omega)}^2 dx ds \right)^{5/9} \quad (4.94)$$

$$\leq C \cdot \|\hat{\Phi}_{tr}\|_{L^5(\Omega_T)}^{10/9} \cdot \left(\|P_1\|_{L^2[(0,T), W^{1,2}(\Omega)]}^{10/9} + \|\partial P_1/\partial s\|_{L^{10/9}[(0,T), (W^{1,2}(\Omega))^*]}^{10/9} \right) \cdot T^{5/9} \quad (4.95)$$

by application of the Aubin-Dubinskij lemma to P_1 . Since $\hat{\Phi}_{tr} \in L^5(\Omega_T)$, we may use (4.55) and (4.88) in order to conclude that

$$\begin{aligned} J_5 &\leq C \left(\left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 + \left\| \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 \right)^{q/2} \\ &\quad + C \left(1 + \left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^{20/9}[(0,T), L^2(\Omega)]}^{20/9} + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^{20/9}[(0,T), L^2(\Omega)]}^{20/9} \right). \end{aligned} \quad (4.96)$$

For the second term, we find

$$J_6 = \sup_{\dots} C \int_0^T \left(\int_{\Omega} |P_3| |\psi| dx \right)^q ds \leq \sup_{\dots} C \int_0^T \left(\|P_3(s)\|_{L^2(\Omega)}^2 + \|\psi(s)\|_{L^2(\Omega)}^2 \right)^{10/9} ds \quad (4.97)$$

$$\leq C \left(1 + \|P_3\|_{L^\infty[(0,T), L^2(\Omega)]}^{20/9} \right) \quad (4.98)$$

$$\leq C \left(1 + \left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 + \left\| \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 \right)^q \quad (4.99)$$

by (4.51). For the third term, we get

$$J_7 = \sup_{\dots} \int_0^T \left(\int_{\Omega} \left| \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right| |\psi| dx \right)^q ds \leq \sup_{\dots} \int_0^T \left(\left\| \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2 \right. \quad (4.100)$$

$$\left. + \|\psi(s)\|_{L^2(\Omega)}^2 \right)^q ds \leq C \left(1 + \left\| \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^{2q}[(0,T), L^2(\Omega)]}^{20/9} \right). \quad (4.101)$$

Combining now (4.91) with (4.96), (4.99) and (4.101), we arrive at the claimed estimate

$$\| \partial P_3 / \partial s \|_{L^q[(0, T), (L^2(\Omega))^*]}^q \quad (4.102)$$

$$\begin{aligned} &\leq C \left(\left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 + \left\| \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 \right)^{q/2} \\ &\quad + C \left(1 + \left\| \frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^{20/9}[(0, T), L^2(\Omega)]}^{20/9} + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^{20/9}[(0, T), L^2(\Omega)]}^{20/9} \right). \end{aligned}$$

• **Step 6.** *The estimate for $\| P_2 \|_{L^2[(0, T), W^{1,2}(\Omega)]}^2$.* Inserting $P_2(s) \in W^{1,2}(\Omega)$ with $\int_{\Omega} P_2(x, s) dx = 0$ as a feasible test function into (4.9), the uniform ellipticity of M_i , M_e and the Poincaré inequality imply

$$\| P_2(s) \|_{W^{1,2}(\Omega)}^2 \leq C \int_{\Omega} \nabla P_2^T (M_i + M_e) \nabla P_2 dx \quad (4.103)$$

$$\leq C \left| \int_{\Omega} \nabla P_2^T M_i \nabla P_1 dx + \int_{\Omega} \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) P_2 dx \right| \quad (4.104)$$

$$\leq C \left(\| P_1 \|_{W^{1,2}(\Omega)} \cdot \| P_2 \|_{W^{1,2}(\Omega)} + \int_{\Omega} \left| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right| | P_2 | dx \right) \quad (4.105)$$

since, by Assumption 2.1., 2), the entries of M_i are essentially bounded. Consequently, applying the generalized Cauchy inequality twice, we get

$$\begin{aligned} \| P_2(s) \|_{W^{1,2}(\Omega)}^2 &\leq C \left(\frac{1}{\varepsilon'_3} \| P_1 \|_{W^{1,2}(\Omega)}^2 + \varepsilon'_3 \| P_2 \|_{W^{1,2}(\Omega)}^2 \right) \\ &\quad + \frac{1}{\varepsilon'_4} \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2 + \varepsilon'_4 \| P_2 \|_{W^{1,2}(\Omega)}^2 \end{aligned} \quad (4.106)$$

for arbitrary $\varepsilon'_3, \varepsilon'_4 > 0$. Choosing $(\varepsilon'_3 + \varepsilon'_4) = 1/(2C)$, we arrive at

$$\frac{1}{2} \| P_2(s) \|_{W^{1,2}(\Omega)}^2 \leq C \left(\| P_1 \|_{W^{1,2}(\Omega)}^2 + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega)}^2 \right) \implies \quad (4.107)$$

$$\| P_2 \|_{L^2[(0, T), W^{1,2}(\Omega)]}^2 \leq C \left(\| P_1 \|_{L^2[(0, T), W^{1,2}(\Omega)]}^2 + \left\| \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \right\|_{L^2(\Omega_T)}^2 \right) \quad (4.108)$$

where the right-hand side is bounded by (4.55).

• **Step 7.** *Conclusion of the proof.* The fact that P_1 belongs even to $C^0[[0, T], L^2(\Omega)]$ can be confirmed analogously to [BOURGAULT/COUDIÈRE/PIERRE 09], p. 478, Subsection 5.3. As a consequence of the imbedding theorem [EVANS 98], p. 286, Theorem 2, $P_3 \in C^0[[0, T], L^2(\Omega)]$ holds true as well. Consequently, the norms on the left-hand side of (4.51) can be replaced by $C^0[[0, T], L^2(\Omega)]$ -norms, and the proof is complete. ■

Proof of Theorem 4.2. • **Step 1.** *Approximate solutions for the reduced adjoint system.* By [BOURGAULT/COUDIÈRE/PIERRE 09], p. 464, Theorem 6, the bidomain bilinear form $A(\cdot, \cdot)$ gives rise to an orthonormal basis of eigenfunctions $\{\psi_i\}$ within the space $W^{1,2}(\Omega)$, which are related to eigenvalues $0 = \lambda_0 < \lambda_1 \leq \lambda_2 \leq \dots$. For $N \in \mathbb{N}_0$, let us define the subspaces

$$X^N(\Omega) = \left\{ \psi = \sum_{i=0}^N c_i \psi_i \mid c_0, \dots, c_N \in \mathbb{R} \right\} \subset W^{1,2}(\Omega), \quad (4.109)$$

and the functions $P_1^N, P_3^N : \Omega \times [0, T] \rightarrow X^N$ according to

$$P_1^N(x, s) = \sum_{i=0}^N p_{i,N}(s) \psi_i(x); \quad P_3^N(x, s) = \sum_{i=0}^N q_{i,N}(s) \psi_i(x) \quad (4.110)$$

where $p_{i,N}, q_{i,N} : [0, T] \rightarrow \mathbb{R}$ are solutions of the initial value problem

$$\begin{aligned} \frac{dp_{j,N}(s)}{ds} + \lambda_j p_{j,N}(s) + \sum_{i=0}^N p_{i,N}(s) \cdot \int_{\Omega} \frac{\partial I_{ion}}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{W}) \psi_i(x) \psi_j(x) dx \\ + \sum_{i=0}^N q_{i,N}(s) \cdot \int_{\Omega} \frac{\partial G}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{W}) \psi_i(x) \psi_j(x) dx = \langle \tilde{S}(s), \psi_j \rangle, \quad 0 \leq j \leq N; \end{aligned} \quad (4.111)$$

$$\begin{aligned} \frac{dq_{j,N}(s)}{ds} + \sum_{i=0}^N p_{i,N}(s) \cdot \int_{\Omega} \frac{\partial I_{ion}}{\partial w}(\hat{\Phi}_{tr}, \hat{W}) \psi_i(x) \psi_j(x) dx \\ + \sum_{i=0}^N q_{i,N}(s) \cdot \int_{\Omega} \frac{\partial G}{\partial w}(\hat{\Phi}_{tr}, \hat{W}) \psi_i(x) \psi_j(x) dx = -\langle \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}), \psi_j \rangle, \quad 0 \leq j \leq N; \end{aligned} \quad (4.112)$$

$$p_{j,N}(0) = 0; \quad q_{j,N}(0) = 0 \quad 0 \leq j \leq N. \quad (4.113)$$

Specifying the data for (4.111) – (4.113) according to the Rogers-McCulloch model, the problem reads as

$$\begin{aligned} \frac{dp_{j,N}(s)}{ds} + \lambda_j p_{j,N}(s) + \sum_{i=0}^N p_{i,N}(s) \cdot \int_{\Omega} \left(3b(\hat{\Phi}_{tr})^2 + 2(a+1)b\hat{\Phi}_{tr} + \hat{W} + ab \right) \psi_i \psi_j dx \\ - \varepsilon \kappa q_{j,N}(s) = \langle \tilde{S}(s), \psi_j \rangle, \quad 0 \leq j \leq N; \end{aligned} \quad (4.114)$$

$$\frac{dq_{j,N}(s)}{ds} + \sum_{i=0}^N p_{i,N}(s) \cdot \int_{\Omega} \hat{\Phi}_{tr} \psi_i \psi_j dx + \varepsilon q_{j,N}(s) = -\langle \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}), \psi_j \rangle, \quad 0 \leq j \leq N; \quad (4.115)$$

$$p_{j,N}(0) = 0; \quad q_{j,N}(0) = 0 \quad 0 \leq j \leq N. \quad (4.116)$$

Obviously, all integrals with respect to x are well-defined and the coefficients as well as the right-hand sides are integrable with respect to s at least. Then, by [WARGA 72], p. 92, Theorem II.4.6., the initial-value problem (4.114) – (4.116) admits a unique solution $(p_{0,N}, \dots, p_{N,N}, q_{0,N}, \dots, q_{N,N}) \in (W^{1,1}(0, T))^{2(N+1)}$. As a consequence of the orthogonality relations, P_1^N and P_3^N obey the equations

$$\begin{aligned} \frac{d}{ds} \langle P_1^N(s), \psi \rangle + A(P_1^N(s), \psi) + \int_{\Omega} \left(\frac{\partial I_{ion}}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{W}) P_1^N + \frac{\partial G}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{W}) P_3^N \right) \psi dx \\ = \langle \tilde{S}(s), \psi \rangle \quad \forall \psi \in X^N(\Omega); \end{aligned} \quad (4.117)$$

$$\begin{aligned} \frac{d}{ds} \langle P_3^N(s), \psi \rangle + \int_{\Omega} \left(\frac{\partial I_{ion}}{\partial w}(\hat{\Phi}_{tr}, \hat{W}) P_1^N + \frac{\partial G}{\partial w}(\hat{\Phi}_{tr}, \hat{W}) P_3^N \right) \psi dx \\ = -\langle \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}), \psi \rangle \quad \forall \psi \in X^N(\Omega). \end{aligned} \quad (4.118)$$

In this sense, the functions P_1^N, P_3^N can be interpreted as approximate solutions of the reduced adjoint system.

• **Step 2.** *A-priori estimates for the approximate solutions P_1^N, P_3^N .* The functions P_1^N, P_3^N obey the a-priori estimates from Theorem 4.1., 2). More precisely, the following holds:

Lemma 4.5. *Let the assumptions of Theorem 4.1., 2), hold for the data of (P) and a feasible solution $(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e)$ where $q = 10/9$. Then for all $N \in \mathbb{N}_0$, the functions P_1^N, P_3^N satisfy the estimate*

$$\begin{aligned} \|P_1^N\|_{C^0[0, T], L^2(\Omega)}^2 + \|P_1^N\|_{L^2((0, T), W^{1,2}(\Omega))}^2 + \|\partial P_1^N / \partial s\|_{L^q((0, T), (W^{1,2}(\Omega))^*)}^q \\ + \|P_3^N\|_{C^0[0, T], L^2(\Omega)}^2 + \|\partial P_3^N / \partial s\|_{L^q((0, T), L^2(\Omega))}^q \leq C \end{aligned} \quad (4.119)$$

for a constant $C > 0$ independent of N .

Proof. We rely on the proof of Theorem 4.1. First, we observe that Lemma 4.3., (4.51) and (4.55) remain true if P_1 and P_3 are replaced by P_1^N and P_3^N since, in Steps 2 and 3 of the proof above, the reduced equations must be studied only for the special test functions $P_1^N(s), P_3^N(s) \in X^N(\Omega)$. Further, we observe that

$$\|\partial P_1^N / \partial s\|_{L^q[(0, T), (W^{1,2}(\Omega))^*]}^q = \int_0^T \sup_{\|\psi\|_{W^{1,2}(\Omega)}=1} |\langle \partial P_1^N(s) / \partial s, \psi \rangle|^q ds \quad (4.120)$$

$$= \int_0^T \sup_{\|\sum_{j=0}^{\infty} c_j \psi_j\|_{W^{1,2}(\Omega)}=1} \left| \left\langle \sum_{i=0}^N \frac{dp_{i,N}(s)}{ds} \psi_i, \sum_{j=0}^{\infty} c_j \psi_j \right\rangle \right|^q ds \quad (4.121)$$

$$= \int_0^T \sup_{\|\sum_{j=0}^{\infty} c_j \psi_j\|_{W^{1,2}(\Omega)}=1} \left| \left\langle \sum_{i=0}^N \frac{dp_{i,N}(s)}{ds} \psi_i, \sum_{j=0}^N c_j \psi_j \right\rangle \right|^q ds \quad (4.122)$$

$$= \int_0^T \sup_{\psi \in X^N, \|\psi\|_{W^{1,2}(\Omega)}=1} |\langle \partial P_1^N(s) / \partial s, \psi \rangle|^q ds. \quad (4.123)$$

By (4.117), the calculations from the proof of Theorem 4.1., Step 4, can be repeated now, resulting in a uniform bound for $\|\partial P_1^N / \partial s\|_{L^q[(0, T), (W^{1,2}(\Omega))^*]}^q$. In the same manner, we may repeat the derivation from Step 5 since

$$\|\partial P_3^N / \partial s\|_{L^q[(0, T), (L^2(\Omega))^*]}^q = \int_0^T \sup_{\|\psi\|_{L^2(\Omega)}=1} |\langle \partial P_3^N(s) / \partial s, \psi \rangle|^q ds \quad (4.124)$$

$$= \int_0^T \sup_{\psi \in X^N, \|\psi\|_{L^2(\Omega)}=1} |\langle \partial P_3^N(s) / \partial s, \psi \rangle|^q ds, \quad (4.125)$$

and we obtain a uniform bound for $\|\partial P_3^N / \partial s\|_{L^q[(0, T), L^2(\Omega)]}^q$ as well. The arguments from Step 7 hold without alterations. ■

• **Step 3.** *The solution for the reduced adjoint system.* Lemma 4.4. implies that we may select a subsequence $\{(P_1^{N'}, P_3^{N'})\}$ of $\{(P_1^N, P_3^N)\}$ with convergence to limit elements in the following sense:

$$P_1^{N'} \rightharpoonup_{L^2[(0, T), W^{1,2}(\Omega)]} P_1; \quad (4.126)$$

$$dP_1^{N'} / ds \rightharpoonup_{L^q[(0, T), (W^{1,2}(\Omega))^*]} \tilde{P}; \quad (4.127)$$

$$P_3^{N'} \rightharpoonup_{L^2[(0, T), L^2(\Omega)]} P_3; \quad (4.128)$$

$$dP_3^{N'} / ds \rightharpoonup_{L^q[(0, T), (L^2(\Omega))^*]} \tilde{Q}. \quad (4.129)$$

Consequently, taking an arbitrary element $\psi_j \in W^{1,2}(\Omega)$ from the orthonormal base, we find

$$\begin{aligned} & \langle \tilde{P}(s), \psi_j \rangle + A(P_1(s), \psi_j) + \int_{\Omega} \left(\frac{\partial I_{ion}}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{W}) P_1 + \frac{\partial G}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{W}) P_3 \right) \psi_j dx - \langle \tilde{S}(s), \psi_j \rangle \\ &= \lim_{N' \rightarrow \infty} \left(\frac{d}{ds} \langle P_1^{N'}(s), \psi_j \rangle + A(P_1^{N'}(s), \psi_j) + \int_{\Omega} \left(\frac{\partial I_{ion}}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{W}) P_1^{N'} \right. \right. \\ & \quad \left. \left. + \frac{\partial G}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{W}) P_3^{N'} \right) \psi_j dx - \langle \tilde{S}(s), \psi_j \rangle \right) = 0 \end{aligned} \quad (4.130)$$

since $\psi_j \in X^{N'}$ for all sufficiently large $N' \in \mathbb{N}$. For the same reason, it holds that

$$\begin{aligned} & \langle \tilde{Q}(s), \psi_j \rangle + \int_{\Omega} \left(\frac{\partial I_{ion}}{\partial w}(\hat{\Phi}_{tr}, \hat{W}) P_1 + \frac{\partial G}{\partial w}(\hat{\Phi}_{tr}, \hat{W}) P_3 \right) \psi_j dx + \left\langle \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}), \psi_j \right\rangle \\ &= \lim_{N' \rightarrow \infty} \left(\frac{d}{ds} \langle P_3^{N'}(s), \psi_j \rangle + \int_{\Omega} \left(\frac{\partial I_{ion}}{\partial w}(\hat{\Phi}_{tr}, \hat{W}) P_1^{N'} + \frac{\partial G}{\partial w}(\hat{\Phi}_{tr}, \hat{W}) P_3^{N'} \right) \psi_j dx \right. \\ & \quad \left. + \left\langle \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}), \psi_j \right\rangle \right) = 0. \end{aligned} \quad (4.131)$$

Weak continuity of the distributional differential operator implies that $\tilde{P} = dP_1/ds$ and $\tilde{Q} = dP_3/ds$ in the sense of distributions. Further, it obviously holds that $P_1(x, 0) = \lim_{N' \rightarrow \infty} P_1^{N'}(x, 0) = 0$ and $P_3(x, 0) = \lim_{N' \rightarrow \infty} P_3^{N'}(x, 0) = 0$. Since $\{\psi_i\}$ lies dense in $W^{1,2}(\Omega)$ as well as in $L^2(\Omega)$, the functions P_1 and P_3 form a weak solution of the reduced adjoint system.

• **Step 4. Completion of the adjoint solution.** As indicated in Subsection 4.2., the solution (P_1, P_3) of the reduced adjoint system may be completed to a weak solution (P_1, P_2, P_3) of the adjoint system where $P_2 \in L^2[(0, T), W^{1,2}(\Omega)]$ with $\int_{\Omega} P_2(x, s) dx = 0$ ($\forall s \in (0, T)$) is uniquely determined by P_1, P_3 . The claimed regularity of the solution is guaranteed by Theorem 4.1., 2).

• **Step 5. Uniqueness.** Since the reduced adjoint system is linear with respect to P_1 and P_3 , estimate (4.20) yields the uniqueness of its weak solution (P_1, P_3) within the space $(L^\infty[(0, T), L^2(\Omega)] \cap L^2[(0, T), W^{1,2}(\Omega)]) \times L^\infty[(0, T), L^2(\Omega)]$. The completion of (P_1, P_3) to a weak solution (P_1, P_2, P_3) of the adjoint system is uniquely determined as well. This finishes the proof of Theorem 4.2. ■

Remarks. 1) If the Rogers-McCulloch model in (3.12) – (3.14) is replaced by the FitzHugh-Nagumo model then the proofs of Theorems 4.1. and 4.2. can be repeated with only minor alterations.

2) Theorems 4.1. and 4.2. remain even true if (3.12) – (3.14) is considered with the linearized Aliev-Panfilov model. In the proofs, we must work with

$$\frac{\partial G}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{W}) = -\varepsilon \kappa (a + 1) + 2\varepsilon \kappa \hat{\Phi}_{tr} \quad (4.132)$$

instead of $\partial G(\hat{\Phi}_{tr}, \hat{W})/\partial \varphi \equiv -\varepsilon \kappa$. Thus the estimations (4.36) – (4.55) have to be modified in the following way: On the right-hand side of (4.36), the term $\varepsilon \kappa \int_{\Omega} |P_1 P_3| dx$ must be replaced by

$$\varepsilon \kappa (a + 1) \int_{\Omega} |P_1 P_3| dx + \varepsilon \kappa \int_{\Omega} |\hat{\Phi}_{tr} P_1 P_3| dx. \quad (4.133)$$

The estimation of the first member of (4.133) runs as above, for the second one we get with arbitrary $\varepsilon_3(s) > 0$:

$$\int_{\Omega} |\hat{\Phi}_{tr} P_1 P_3| dx \leq C \varepsilon_3(s) \int_{\Omega} |\hat{\Phi}_{tr} P_1|^2 dx + \frac{C}{\varepsilon_3(s)} \|P_3\|_{L^2(\Omega)}^2 \quad (4.134)$$

$$\leq C \varepsilon_3(s) \|\hat{\Phi}_{tr}\|_{L^4(\Omega)}^2 \cdot \|P_1\|_{L^4(\Omega)}^2 + \frac{C}{\varepsilon_3(s)} \|P_3\|_{L^2(\Omega)}^2. \quad (4.135)$$

We choose $\varepsilon_3(s) = \varepsilon'_3 / (1 + \|\hat{\Phi}_{tr}\|_{L^4(\Omega)}^2)$ with $\varepsilon'_3 > 0$, thus getting

$$\int_{\Omega} |\hat{\Phi}_{tr} P_1 P_3| dx \leq C \varepsilon' \|P_1\|_{W^{1,2}(\Omega)}^2 + \frac{C}{\varepsilon'_3} (1 + \|\hat{\Phi}_{tr}\|_{L^4(\Omega)}^2) \|P_3\|_{L^2(\Omega)}^2, \quad (4.136)$$

and with appropriate choices of $\varepsilon'_3 > 0$, we may proceed as above. Further alterations concern the estimations (4.58) – (4.88). In (4.58), the term $\sup \dots \varepsilon^q \kappa^q \left(\int_{\Omega} |P_3| |\psi| dx \right)^q$ must be replaced by

$$\sup \dots \varepsilon^q \kappa^q (a+1)^q \left(\int_{\Omega} |P_3| |\psi| dx \right)^q + \sup \dots \varepsilon^q \kappa^q \left(\int_{\Omega} |\hat{\Phi}_{tr} P_3 \psi| dx \right)^q. \quad (4.137)$$

Despite of the lesser regularity of \hat{W} for the linearized Aliev-Panfilov model, the estimations (4.70) – (4.80) can be maintained since the solution satisfies $\hat{W} \in C^0[[0, T], L^{8/3}(\Omega)] \hookrightarrow L^{10/4}[(0, T), L^{20/11}(\Omega)]$. In (4.81) – (4.84), we must add an estimate for the second term from (4.137). Consider therefore

$$\begin{aligned} & \sup \dots \int_0^T \int_{\Omega} |\hat{\Phi}_{tr}|^{10/9} |P_3|^{10/9} |\psi|^{10/9} dx ds \\ & \leq C \sup \dots \int_0^T \left(\int_{\Omega} |\hat{\Phi}_{tr}|^5 dx \right)^{4/18} \left(\int_{\Omega} |\psi|^5 dx \right)^{4/18} \left(\int_{\Omega} |P_3|^2 dx \right)^{10/18} ds \end{aligned} \quad (4.138)$$

$$\leq C \sup \dots \int_0^T \left(\|\hat{\Phi}_{tr}\|_{L^5(\Omega)}^{10/9} \cdot \|\psi\|_{W^{1,2}(\Omega)}^{10/9} \cdot \|P_3\|_{L^2(\Omega)}^{10/9} \right) ds = C \int_0^T \left(\|\hat{\Phi}_{tr}\|_{L^5(\Omega)}^{10/9} \cdot \|P_3\|_{L^2(\Omega)}^{10/9} \right) ds \quad (4.139)$$

$$\leq C \|\hat{\Phi}_{tr}\|_{L^{10/9}[(0, T), L^5(\Omega)]}^{10/9} \cdot \|P_3\|_{L^{\infty}[(0, T), L^2(\Omega)]}^{10/9}, \quad (4.140)$$

and we may proceed as above. The remaining parts of the proof of Theorem 4.1. as well as the proof of Theorem 4.2. remain unchanged.

5. Necessary optimality conditions.

5.1. Statement of the theorems.

Definition 5.1. (Weak local minimizer) A quadruple $(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e)$, which is feasible in (P), is called a weak local minimizer of (P) iff there exists a number $\varepsilon > 0$ such that for all admissible $(\Phi_{tr}, \Phi_e, W, I_e)$ the conditions

$$\|\Phi_{tr} - \hat{\Phi}_{tr}\|_{X_1} \leq \varepsilon, \quad \|\Phi_e - \hat{\Phi}_e\|_{X_2} \leq \varepsilon, \quad \|W - \hat{W}\|_{X_3} \leq \varepsilon, \quad \|I_e - \hat{I}_e\|_{X_4} \leq \varepsilon \quad (5.1)$$

imply the relation $F(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e) \leq F(\Phi_{tr}, \Phi_e, W, I_e)$.

The necessary optimality conditions for weak local minimizers of (P) can be formulated as follows:

Theorem 5.2. (First-order necessary optimality conditions for the control problem (P)) We consider problem (P), (3.11) – (3.17), under the assumptions of Subsection 3.1. with the Rogers-McCulloch or the FitzHugh-Nagumo model. Assume further that 1) $\Omega \subset \mathbb{R}^3$ admits a $C^{1,1}$ -boundary, 2) $M_i, M_e: \text{cl}(\Omega) \rightarrow \mathbb{R}^{3 \times 3}$ are symmetric, positive definite matrix functions obeying (2.5) with $W^{1,\infty}(\Omega)$ -coefficients, and 3) the integrand $r(x, t, \varphi, \eta, w)$ is continuously differentiable with respect to φ, η and w . Let $(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e)$ be a weak local minimizer of (P) such that $(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W})$ is a strong solution of the bidomain system on $[0, T]$, $\hat{I}_e \in W^{1,2}[(0, T), L^2(\Omega)]$ and

$$\frac{\partial r}{\partial \varphi}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}), \quad \frac{\partial r}{\partial \eta}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}), \quad \frac{\partial r}{\partial w}(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) \in L^{2q}[(0, T), L^2(\Omega)] \quad (5.2)$$

where $q \geq 10/9$. Then there exist multipliers $P_1 \in L^4[(0, T), W^{1,2}(\Omega)]$, $P_2 \in L^2[(0, T), W^{1,2}(\Omega)] \cap \{Z \mid \int_{\Omega} Z(x, t) dx = 0 \ (\forall) t \in (0, T)\}$ and $P_3 \in L^2(\Omega_T)$, satisfying together with $(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e)$ the adjoint equations (4.8) – (4.10), which are solved in weak sense, as well as the optimality condition

$$\int_0^T \int_{\Omega_{con}} \left(\mu \hat{I}_e - Q P_2 \right) \cdot (I_e - \hat{I}_e) dx dt \geq 0 \quad \forall I_e \in C. \quad (5.3)$$

If the linearized Aliev-Panfilov model is specified then all assertions remain true provided that $\partial r(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W})/\partial w$ belongs to $L^3(\Omega_T)$ instead of $L^2(\Omega_T)$.

The assumptions of Theorem 5.2. reflect the fact that there is a regularity gap between the weak solutions of the primal and adjoint equations. The duality pairing between $\partial\Phi_{tr}/\partial t \in L^{4/3}[(0, T), (W^{1,2}(\Omega))^*]$ and $P_1 \in L^2[(0, T), W^{1,2}(\Omega)]$ is not well-defined, and hence further regularity is required. In order to gain this regularity, we have to impose that $(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W})$ is a *strong* solution of the bidomain system rather than a weak one. Sufficient conditions for strong local solvability of (3.12) – (3.14) may be found in [VENERONI 09].

Corollary 5.3. (Pointwise formulation of the optimality condition) *Under the assumptions of Theorem 5.2., let the optimal control \hat{I}_e be represented as $\hat{I}_e = Q \hat{I}$ with $\hat{I} \in L^\infty[(0, T), L^2(\Omega)]$, $\text{supp}(\hat{I}) \subseteq \Omega_{con} \times [0, T]$ and $|\hat{I}(x, t)| \leq R$ for almost all $(x, t) \in \Omega_T$. The optimality condition (5.3) from Theorem 5.2. then implies the following Pontryagin minimum condition, which holds a. e. pointwise:*

$$\hat{I}(x_0, t_0) \cdot \left(\mu \cdot Q \hat{I}(x_0, t_0) - Q P_2(x_0, t_0) \right) = \underset{-R \leq \eta \leq R}{\text{Min}} \eta \left(\mu \cdot Q \hat{I}(x_0, t_0) - Q P_2(x_0, t_0) \right) \quad (5.4)$$

$$(\forall) (x_0, t_0) \in \Omega_{con} \times [0, T].$$

Consequently, for a. e. $(x, t) \in \Omega_{con} \times [0, T]$ the following implications hold:

$$Q \hat{I}(x, t) - \frac{1}{\mu} Q P_2(x, t) > 0 \implies \hat{I}(x, t) = -R;$$

$$Q \hat{I}(x, t) - \frac{1}{\mu} Q P_2(x, t) < 0 \implies \hat{I}(x, t) = R \text{ and} \quad (5.5)$$

$$\hat{I}(x, t) \in (-R, R) \implies Q \hat{I}(x, t) - \frac{1}{\mu} Q P_2(x, t) = 0.$$

Corollary 5.4. (Regularity of weak local minimizers) *Under the assumptions of Theorem 5.2., consider a weak local minimizer $(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e)$ of (P), whose control part $\hat{I}_e = Q \hat{I}$ is generated by a function \hat{I} with $|\hat{I}(x, t)| < R$ a. e. Then $\hat{I}_e|_{\Omega_{con}}$ belongs to the space $L^\infty(\Omega_{con} \times [0, T]) \cap L^2[(0, T), W^{1,2}(\Omega_{con})]$.*

For numerical purposes, it is useful to specify the Gâteaux derivative of the reduced cost functional $\tilde{F}: \mathcal{C} \rightarrow \mathbb{R}$. It is defined through

$$\tilde{F}(I_e) = F(\Phi_{tr}(I_e), \Phi_e(I_e), W(I_e), I_e) \quad (5.6)$$

with the aid of the control-to-state mapping $I_e \mapsto (\Phi_{tr}(I_e), \Phi_e(I_e), W(I_e))$, which is well-defined by Proposition 3.1.

Corollary 5.5. (First variation of the reduced cost functional) *Under the assumptions of Theorem 5.2., the Gâteaux derivative of the reduced cost functional \tilde{F} at $\hat{I}_e \in \mathcal{C}$ is given through*

$$D_{I_e} \tilde{F}(\hat{I}_e) = \mu \hat{I}_e - Q P_2(\hat{I}_e) \quad (5.7)$$

where $(P_1(\hat{I}_e), P_2(\hat{I}_e), P_3(\hat{I}_e))$ denotes the solution of the adjoint system (4.8) – (4.10) corresponding to $(\Phi_{tr}(\hat{I}_e), \Phi_e(\hat{I}_e), W(\hat{I}_e), \hat{I}_e)$.

5.2. Proof of the necessary optimality conditions.

Proof of Theorem 5.2. As mentioned in the introduction, the proof of the necessary optimality conditions for (P) is based on the stability estimate for the bidomain system (Theorem 2.4.) and the existence theorem for the adjoint system (Theorem 4.2.), which will be invoked in Steps 2 and 3 of the proof, respectively.

• **Step 1.** *Variation of the weak local minimizer in a feasible direction.* Assume that $(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e)$ is a weak local minimizer of (P). If $I_e \in \mathcal{C}$ is an arbitrary feasible control with $\|I_e - \hat{I}_e\|_{L^\infty[(0, T), L^2(\Omega)]} \leq \varepsilon$ then, by Proposition 3.2., all controls

$$I_e(s) = \hat{I}_e + s(I_e - \hat{I}_e), \quad 0 \leq s \leq 1, \quad (5.8)$$

belong to \mathcal{C} as well. By Proposition 3.1., for every $I_e(s) \in L^\infty[(0, T), L^2(\Omega)]$, there exists a corresponding weak solution $(\Phi_{tr}(s), \Phi_e(s), W(s)) \in X_1 \times X_2 \times X_3$ for the bidomain system on $[0, T]$. Thus the quadruples $(\Phi_{tr}(s), \Phi_e(s), W(s), I_e(s))$ are feasible in (P) for all $0 \leq s \leq 1$. On the other hand, from [KUNISCH/WAGNER 11], p. 7, Theorem 2.7., it follows that every feasible solution of (P) within a closed ball

$$U_\varepsilon(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e) = K(\hat{\Phi}_{tr}, C\varepsilon) \times K(\hat{\Phi}_e, C\varepsilon) \times K(\hat{W}, C\varepsilon) \times K(\hat{I}_e, \varepsilon) \subset X_1 \times X_2 \times X_3 \times X_4 \quad (5.9)$$

can be generated in this way.

• **Step 2. Lemma 5.6.** *For all $I_e \in \mathcal{C}$, $\|I_e - \hat{I}_e\|_{L^\infty[(0, T), L^2(\Omega)]} \leq \varepsilon$ implies that*

$$\lim_{s \rightarrow 0+0} \frac{1}{s} \|\Phi_{tr}(s) - \hat{\Phi}_{tr}\|_{X_1}^2 = 0; \quad \lim_{s \rightarrow 0+0} \frac{1}{s} \|\Phi_{tr}(s) - \hat{\Phi}_{tr}\|_{\tilde{X}_1}^2 = 0; \quad (5.10)$$

$$\lim_{s \rightarrow 0+0} \frac{1}{s} \|\Phi_e(s) - \hat{\Phi}_e\|_{X_2}^2 = 0; \quad (5.11)$$

$$\lim_{s \rightarrow 0+0} \frac{1}{s} \|W(s) - \hat{W}\|_{X_3}^2 = 0 \quad \text{and} \quad \lim_{s \rightarrow 0+0} \frac{1}{s} \|W(s) - \hat{W}\|_{\tilde{X}_3}^2 = 0. \quad (5.12)$$

Proof. The stability estimate [KUNISCH/WAGNER 11], p. 7, Theorem 2.7., (2.38), implies

$$\begin{aligned} \|\Phi_{tr}(s) - \hat{\Phi}_{tr}\|_{X_1}^2 &= \|\Phi_{tr}(s) - \hat{\Phi}_{tr}\|_{L^2[(0, T), W^{1,2}(\Omega)]}^2 \leq C \cdot \|I_e(s) - \hat{I}_e\|_{L^\infty[(0, T), (W^{1,2}(\Omega))^*]}^2 \\ &\leq C \cdot \|I_e(s) - \hat{I}_e\|_{L^\infty[(0, T), L^2(\Omega)]}^2 = C s^2 \|I_e - \hat{I}_e\|_{L^\infty[(0, T), L^2(\Omega)]}^2 \implies \end{aligned} \quad (5.13)$$

$$\lim_{s \rightarrow 0+0} \frac{1}{s} \|\Phi_{tr}(s) - \hat{\Phi}_{tr}\|_{X_1}^2 \leq \lim_{s \rightarrow 0+0} C s \|I_e - \hat{I}_e\|_{L^\infty[(0, T), L^2(\Omega)]}^2 = 0 \quad (5.14)$$

as well as

$$\begin{aligned} \|\Phi_e(s) - \hat{\Phi}_e\|_{X_2}^2 &= \|\Phi_e(s) - \hat{\Phi}_e\|_{L^2[(0, T), W^{1,2}(\Omega)]}^2 \leq C \cdot \|I_e(s) - \hat{I}_e\|_{L^\infty[(0, T), (W^{1,2}(\Omega))^*]}^2 \\ &\leq C \cdot \|I_e(s) - \hat{I}_e\|_{L^\infty[(0, T), L^2(\Omega)]}^2 = C s^2 \|I_e - \hat{I}_e\|_{L^\infty[(0, T), L^2(\Omega)]}^2 \implies \end{aligned} \quad (5.15)$$

$$\lim_{s \rightarrow 0+0} \frac{1}{s} \|\Phi_e(s) - \hat{\Phi}_e\|_{X_2}^2 \leq \lim_{s \rightarrow 0+0} C s \|I_e - \hat{I}_e\|_{L^\infty[(0, T), L^2(\Omega)]}^2 = 0 \quad (5.16)$$

and

$$\begin{aligned} \|W(s) - \hat{W}\|_{X_3}^2 &= \|W(s) - \hat{W}\|_{L^2(\Omega_T)}^2 \leq C \cdot \|I_e(s) - \hat{I}_e\|_{L^\infty[(0, T), (W^{1,2}(\Omega))^*]}^2 \\ &\leq C \cdot \|I_e(s) - \hat{I}_e\|_{L^\infty[(0, T), L^2(\Omega)]}^2 = C s^2 \|I_e - \hat{I}_e\|_{L^\infty[(0, T), L^2(\Omega)]}^2 \implies \end{aligned} \quad (5.17)$$

$$\lim_{s \rightarrow 0+0} \frac{1}{s} \|W(s) - \hat{W}\|_{X_3}^2 \leq \lim_{s \rightarrow 0+0} C s \|I_e - \hat{I}_e\|_{L^\infty[(0, T), L^2(\Omega)]}^2 = 0. \quad (5.18)$$

In an analogous manner, the relation with $\|W(s) - \hat{W}\|_{\tilde{X}_3}^2$ can be confirmed. In order to establish the relation with $\|\Phi_{tr}(s) - \hat{\Phi}_{tr}\|_{\tilde{X}_1}^2$, we rely on [KUNISCH/WAGNER 11] p. 7, Theorem 2.7., (2.39), which leads to

$$\|\Phi_{tr}(s) - \hat{\Phi}_{tr}\|_{\tilde{X}_1}^2 = \|\Phi_{tr}(s) - \hat{\Phi}_{tr}\|_{W^{1,4/3}[(0,T), (W^{1,2}(\Omega))^*]}^2 \quad (5.19)$$

$$\leq C^2 \cdot \text{Max} \left(\|I_e(s) - \hat{I}_e\|_{L^\infty[(0,T), (W^{1,2}(\Omega))^*]}^2, \|I_e(s) - \hat{I}_e\|_{L^\infty[(0,T), (W^{1,2}(\Omega))^*]}^4 \right) \quad (5.20)$$

$$\leq C \cdot \text{Max} \left(s^2 \|I_e - \hat{I}_e\|_{L^\infty[(0,T), L^2(\Omega)]}^2, s^4 \|I_e - \hat{I}_e\|_{L^\infty[(0,T), L^2(\Omega)]}^4 \right) \implies \quad (5.21)$$

$$\lim_{s \rightarrow 0+0} \frac{1}{s} \|\Phi_{tr}(s) - \hat{\Phi}_{tr}\|_{\tilde{X}_1}^2 \quad (5.22)$$

$$\leq \lim_{s \rightarrow 0+0} C \cdot \text{Max} \left(s \|I_e - \hat{I}_e\|_{L^\infty[(0,T), L^2(\Omega)]}^2, s^3 \|I_e - \hat{I}_e\|_{L^\infty[(0,T), L^2(\Omega)]}^4 \right) = 0. \quad \blacksquare$$

• **Step 3.** By Theorems 4.1. and 4.2., in correspondence to $(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e)$, there exist functions $P_1 \in (L^{4/3}[(0,T), (W^{1,2}(\Omega))^*])^* = L^4[(0,T), W^{1,2}(\Omega)]$, $P_2 \in (L^2[(0,T), W^{1,2}(\Omega)])^* = L^2[(0,T), W^{1,2}(\Omega)]$ with $\int_\Omega P_2(x,t) dx = 0$ for almost all $t \in (0, T)$ and $P_3 \in (L^2[(0,T), (L^2(\Omega))^*])^* = L^2(\Omega_T)$ satisfying the system (4.8) – (4.10) as weak solutions. Consequently, P_1, P_2 and P_3 solve the adjoint equations (4.5) – (4.7) together with $P_4 = -P_1(\cdot, 0)$ and $P_5 = -P_3(\cdot, 0)$. With these functions, we may derive the following estimates:

Lemma 5.7. *The following estimates hold true:*

$$\lim_{s \rightarrow 0+0} \frac{1}{s} \langle P_1, D_{\hat{\Phi}_{tr}} E_1(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W})(\Phi_{tr}(s) - \hat{\Phi}_{tr}) + D_{\hat{\Phi}_e} E_1(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W})(\Phi_e(s) - \hat{\Phi}_e) \quad (5.23)$$

$$+ D_W E_1(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W})(W(s) - \hat{W}) \rangle = 0;$$

$$\lim_{s \rightarrow 0+0} \frac{1}{s} \langle P_2, D_{\hat{\Phi}_{tr}} E_2(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{I}_e)(\Phi_{tr}(s) - \hat{\Phi}_{tr}) + D_{\hat{\Phi}_e} E_2(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{I}_e)(\Phi_e(s) - \hat{\Phi}_e) \quad (5.24)$$

$$+ \langle P_2, D_{I_e} E_2(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{I}_e)(I_e - \hat{I}_e) \rangle = 0;$$

$$\lim_{s \rightarrow 0+0} \frac{1}{s} \langle P_3, D_{\hat{\Phi}_{tr}} E_3(\hat{\Phi}_{tr}, \hat{W})(\Phi_{tr}(s) - \hat{\Phi}_{tr}) + D_W E_3(\hat{\Phi}_{tr}, \hat{W})(W(s) - \hat{W}) \rangle = 0. \quad (5.25)$$

Proof. We restrict ourselves to the proof of (5.23), noting that (5.24) and (5.25) can be confirmed in a completely analogous manner. Due to our assumptions on the differentiability of r , the principal theorem of calculus in its Bochner integral version is applicable, cf. [BERGER 77], p. 68, (2.1.11). For the feasible solutions $(\Phi_{tr}(s), \Phi_e(s), W(s), I_e(s))$ and $(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e)$, we get from the first state equation in (P), (3.12):

$$\begin{aligned} \mathfrak{o} &= E_1(\Phi_{tr}(s), \Phi_e(s), W(s)) - E_1(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) = \int_0^1 D_{(\Phi_{tr}, \Phi_e, W)} E_1(\hat{\Phi}_{tr} + \tau(\Phi_{tr}(s) - \hat{\Phi}_{tr}), \\ &\quad \hat{\Phi}_e + \tau(\Phi_e(s) - \hat{\Phi}_e), \hat{W} + \tau(W(s) - \hat{W})) (\Phi_{tr}(s) - \hat{\Phi}_{tr}, \Phi_e(s) - \hat{\Phi}_e, W(s) - \hat{W}) d\tau \implies \end{aligned} \quad (5.26)$$

$$\begin{aligned} 0 &= \langle P_1, \int_0^1 \left(D_{(\Phi_{tr}, \Phi_e, W)} E_1(\hat{\Phi}_{tr} + \tau(\Phi_{tr}(s) - \hat{\Phi}_{tr}), \hat{\Phi}_e + \tau(\Phi_e(s) - \hat{\Phi}_e), \hat{W} + \tau(W(s) - \hat{W})) \right. \\ &\quad \left. (\Phi_{tr}(s) - \hat{\Phi}_{tr}, \Phi_e(s) - \hat{\Phi}_e, W(s) - \hat{W}) \right. \\ &\quad \left. - D_{(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W})} E_1(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W})(\Phi_{tr}(s) - \hat{\Phi}_{tr}, \Phi_e(s) - \hat{\Phi}_e, W(s) - \hat{W}) \right) d\tau \rangle \\ &+ \langle P_1, D_{(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W})} E_1(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W})(\Phi_{tr}(s) - \hat{\Phi}_{tr}, \Phi_e(s) - \hat{\Phi}_e, W(s) - \hat{W}) \rangle \end{aligned} \quad (5.27)$$

$$\begin{aligned} &= \langle P_1, \int_0^1 \left(D_{\hat{\Phi}_{tr}} E_1(\hat{\Phi}_{tr} + \tau \dots, \hat{\Phi}_e + \tau \dots, \hat{W} + \tau \dots)(\Phi_{tr}(s) - \hat{\Phi}_{tr}) - D_{\hat{\Phi}_{tr}} E_1(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W})(\Phi_{tr}(s) - \hat{\Phi}_{tr}) \right. \\ &\quad \left. + D_{\hat{\Phi}_e} E_1(\hat{\Phi}_{tr} + \tau \dots, \hat{\Phi}_e + \tau \dots, \hat{W} + \tau \dots)(\Phi_e(s) - \hat{\Phi}_e) - D_{\hat{\Phi}_e} E_1(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W})(\Phi_e(s) - \hat{\Phi}_e) \right. \\ &\quad \left. + D_W E_1(\hat{\Phi}_{tr} + \tau \dots, \hat{\Phi}_e + \tau \dots, \hat{W} + \tau \dots)(W(s) - \hat{W}) - D_W E_1(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W})(W(s) - \hat{W}) \right) d\tau \rangle \end{aligned} \quad (5.28)$$

$$+ \langle P_1, D_{\Phi_{tr}} E_1(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W})(\Phi_{tr}(s) - \hat{\Phi}_{tr}) + D_{\Phi_e} E_1(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W})(\Phi_e(s) - \hat{\Phi}_e) + D_W E_1(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W})(W(s) - \hat{W}) \rangle.$$

By [YOSIDA 95], p. 133, Corollary 1, we have

$$|\langle P_1, \int_0^1 (\dots) d\tau \rangle| \leq \|P_1\|_{Z_1^*} \cdot \left\| \int_0^1 (\dots) d\tau \right\|_{Z_1} \leq \|P_1\|_{Z_1^*} \cdot \int_0^1 \|\dots\|_{Z_1} d\tau. \quad (5.29)$$

Consequently, for the first summand within (5.28), it holds that

$$\begin{aligned} \lim_{s \rightarrow 0+0} \frac{1}{s} |\langle P_1, \int_0^1 (\dots) d\tau \rangle| &\leq \lim_{s \rightarrow 0+0} \|P_1\|_{Z_1^*} \left(\int_0^1 \|D_{\Phi_{tr}} E_1(\hat{\Phi}_{tr} + \tau \dots, \hat{\Phi}_e + \tau \dots, \hat{W} + \tau \dots) \right. \\ &\quad \left. - D_{\Phi_{tr}} E_1(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W})\|_{\mathcal{L}(\tilde{X}_1, Z_1)} \frac{1}{s} \|\Phi_{tr}(s) - \hat{\Phi}_{tr}\|_{\tilde{X}_1} d\tau \right. \\ &\quad \left. + \int_0^1 \|D_{\Phi_e} E_1(\hat{\Phi}_{tr} + \tau \dots, \hat{\Phi}_e + \tau \dots, \hat{W} + \tau \dots) - D_{\Phi_e} E_1(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W})\|_{\mathcal{L}(X_2, Z_1)} \frac{1}{s} \|\Phi_e(s) - \hat{\Phi}_e\|_{X_2} d\tau \right. \\ &\quad \left. + \int_0^1 \|D_W E_1(\hat{\Phi}_{tr} + \tau \dots, \hat{\Phi}_e + \tau \dots, \hat{W} + \tau \dots) - D_W E_1(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W})\|_{\mathcal{L}(\tilde{X}_3, Z_1)} \frac{1}{s} \|W(s) - \hat{W}\|_{\tilde{X}_3} d\tau \right) \\ &\leq \lim_{s \rightarrow 0+0} \|P_1\|_{Z_1^*} \left(\int_0^1 L_1 \tau \left(\|\Phi_{tr}(s) - \hat{\Phi}_{tr}\|_{\tilde{X}_1} + \|\Phi_e(s) - \hat{\Phi}_e\|_{X_2} + \|W(s) - \hat{W}\|_{\tilde{X}_3} \right) \frac{1}{s} \|\Phi_{tr}(s) - \hat{\Phi}_{tr}\|_{\tilde{X}_1} d\tau \right. \\ &\quad \left. + \int_0^1 L_2 \tau \left(\|\Phi_{tr}(s) - \hat{\Phi}_{tr}\|_{\tilde{X}_1} + \|\Phi_e(s) - \hat{\Phi}_e\|_{X_2} + \|W(s) - \hat{W}\|_{\tilde{X}_3} \right) \frac{1}{s} \|\Phi_e(s) - \hat{\Phi}_e\|_{X_2} d\tau \right. \\ &\quad \left. + \int_0^1 L_3 \tau \left(\|\Phi_{tr}(s) - \hat{\Phi}_{tr}\|_{\tilde{X}_1} + \|\Phi_e(s) - \hat{\Phi}_e\|_{X_2} + \|W(s) - \hat{W}\|_{\tilde{X}_3} \right) \frac{1}{s} \|W(s) - \hat{W}\|_{\tilde{X}_3} d\tau \right) \quad (5.31) \end{aligned}$$

with Lipschitz constants L_1, L_2, L_3 , whose existence is ensured by the twice continuous Fréchet differentiability of E_1 with respect to Φ_{tr}, Φ_e and W . With reference to Lemma 5.6., the estimate (5.31) may be continued as follows:

$$\begin{aligned} \lim_{s \rightarrow 0+0} \frac{1}{s} |\langle P_1, \int_0^1 (\dots) d\tau \rangle| &\leq \lim_{s \rightarrow 0+0} \|P_1\|_{Z_1^*} \frac{1}{2} (L_1 + L_2 + L_3) \frac{1}{s} \left(\|\Phi_{tr}(s) - \hat{\Phi}_{tr}\|_{\tilde{X}_1} + \|\Phi_e(s) - \hat{\Phi}_e\|_{X_2} + \|W(s) - \hat{W}\|_{\tilde{X}_3} \right)^2 \quad (5.32) \\ &\leq \lim_{s \rightarrow 0+0} \|P_1\|_{Z_1^*} C \left(\frac{1}{s} \|\Phi_{tr}(s) - \hat{\Phi}_{tr}\|_{\tilde{X}_1}^2 + \frac{1}{s} \|\Phi_e(s) - \hat{\Phi}_e\|_{X_2}^2 + \frac{1}{s} \|W(s) - \hat{W}\|_{\tilde{X}_3}^2 \right) = 0, \quad (5.33) \end{aligned}$$

and this implies the first of the claimed relations, namely

$$\begin{aligned} \lim_{s \rightarrow 0+0} \frac{1}{s} \langle P_1, D_{\Phi_{tr}} E_1(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W})(\Phi_{tr}(s) - \hat{\Phi}_{tr}) + D_{\Phi_e} E_1(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W})(\Phi_e(s) - \hat{\Phi}_e) \\ + D_W E_1(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W})(W(s) - \hat{W}) \rangle = 0. \quad (5.34) \end{aligned}$$

From the second and third state equations (3.13) and (3.14), the limit relations (5.24) and (5.25) can be derived in a completely analogous way. ■

Since $\Phi_{tr}(s)$ and $W(s)$ take the same initial values as $\hat{\Phi}_{tr}$ and \hat{W} , respectively, it holds further that

$$\lim_{s \rightarrow 0+0} \frac{1}{s} \langle P_4, D_{\Phi_{tr}} E_4(\hat{\Phi}_{tr})(\Phi_{tr}(s) - \hat{\Phi}_{tr}) \rangle = \lim_{s \rightarrow 0+0} \frac{1}{s} \langle P_5, D_W E_5(\hat{W})(W(s) - \hat{W}) \rangle = 0. \quad (5.35)$$

• **Step 4.** *The first variation of the objective.* Choose now $\varepsilon > 0$ small enough in order to ensure that the difference $F(\Phi_{tr}(s), \Phi_e(s), W(s), I_e(s)) - F(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e)$ of the objective values is nonnegative for all

quadruples $(\Phi_{tr}(s), \Phi_e(s), W(s), I_e(s))$ belonging to the closed ball $U_\varepsilon(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e)$ defined in (5.9). As a consequence of our assumptions about the integrand r , the first variation may be written as

$$0 \leq \delta^+ F(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e) (\Phi_{tr}(1) - \hat{\Phi}_{tr}, \Phi_e(1) - \hat{\Phi}_e, W(1) - \hat{W}, I_e - \hat{I}_e) \\ = \lim_{s \rightarrow 0+0} \frac{1}{s} \left(F(\Phi_{tr}(s), \Phi_e(s), W(s), I_e(s)) - F(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e) \right) \quad (5.36)$$

$$= \lim_{s \rightarrow 0+0} \frac{1}{s} \left(D_{\Phi_{tr}} F(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e) (\Phi_{tr}(s) - \hat{\Phi}_{tr}) + D_{\Phi_e} F(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e) (\Phi_e(s) - \hat{\Phi}_e) \right. \\ \left. + D_W F(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e) (W(s) - \hat{W}) + D_{I_e} F(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e) (I_e(s) - \hat{I}_e) \right). \quad (5.37)$$

Together with Lemma 5.7. and (5.35), we obtain

$$0 \leq \lim_{s \rightarrow 0+0} \frac{1}{s} \left(D_{\Phi_{tr}} F(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e) (\Phi_{tr}(s) - \hat{\Phi}_{tr}) \right. \\ \left. + \langle P_1, D_{\Phi_{tr}} E_1(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) (\Phi_{tr}(s) - \hat{\Phi}_{tr}) \rangle + \langle P_2, D_{\Phi_{tr}} E_2(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{I}_e) (\Phi_{tr}(s) - \hat{\Phi}_{tr}) \rangle \right. \\ \left. + \langle P_3, D_{\Phi_{tr}} E_3(\hat{\Phi}_{tr}, \hat{W}) (\Phi_{tr}(s) - \hat{\Phi}_{tr}) \rangle + \langle P_4, D_{\Phi_{tr}} E_4(\hat{\Phi}_{tr}) (\Phi_{tr}(s) - \hat{\Phi}_{tr}) \rangle \right. \\ \left. + D_{\Phi_e} F(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e) (\Phi_e(s) - \hat{\Phi}_e) \right. \\ \left. + \langle P_1, D_{\Phi_e} E_1(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) (\Phi_e(s) - \hat{\Phi}_e) \rangle + \langle P_2, D_{\Phi_e} E_2(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{I}_e) (\Phi_e(s) - \hat{\Phi}_e) \rangle \right. \\ \left. + D_W F(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e) (W(s) - \hat{W}) \right. \\ \left. + \langle P_1, D_W E_1(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}) (W(s) - \hat{W}) \rangle + \langle P_2, D_W E_2(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{I}_e) (W(s) - \hat{W}) \rangle \right. \\ \left. + \langle P_3, D_W E_3(\hat{\Phi}_{tr}, \hat{W}) (W(s) - \hat{W}) \rangle + \langle P_5, D_W E_5(\hat{W}) (W(s) - \hat{W}) \rangle \right. \\ \left. + D_{I_e} F(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e) (I_e(s) - \hat{I}_e) + \langle P_2, D_{I_e} E_2(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{I}_e) (I_e - \hat{I}_e) \rangle \right) \quad (5.38)$$

where the first three parts vanish since P_1, P_2, P_3 together with $P_4 = -P_1(\cdot, 0)$ and $P_5 = -P_3(\cdot, 0)$ solve the adjoint equations (4.5) – (4.7). Note that, by Subsection 4.1. above, these equations take the claimed form. Consequently, we arrive at

$$0 \leq \lim_{s \rightarrow 0+0} \frac{1}{s} \left(D_{I_e} F(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{W}, \hat{I}_e) (I_e(s) - \hat{I}_e) + \langle P_2, D_{I_e} E_2(\hat{\Phi}_{tr}, \hat{\Phi}_e, \hat{I}_e) (I_e - \hat{I}_e) \rangle \right) \quad (5.39)$$

$$= \int_0^T \int_\Omega (\mu \hat{I}_e - P_2) \cdot (I_e - \hat{I}_e) dx dt = \int_0^T \int_\Omega (\mu \hat{I}_e - Q P_2) \cdot (I_e - \hat{I}_e) dx dt \quad (5.40)$$

for arbitrary $I_e \in \mathcal{C}$. Since I_e and \hat{I}_e vanish outside $\Omega_{con} \times [0, T]$, this confirms the claimed optimality condition (5.3), and the proof is complete. ■

Proof of Corollary 5.3. Using the representations $I_e = QI$ and $\hat{I}_e = Q\hat{I}$, inequality (5.40) may be rewritten as

$$0 \leq \int_0^T \int_\Omega (\mu \cdot Q\hat{I} - Q P_2) \cdot (QI - Q\hat{I}) dx dt = \int_0^T \int_\Omega (\mu \cdot Q\hat{I} - Q P_2) \cdot (I - \hat{I}) dx dt \quad (5.41)$$

$$= \int_0^T \int_{\Omega_{con}} (\mu \cdot Q\hat{I} - Q P_2) \cdot (I - \hat{I}) dx dt \quad (5.42)$$

$$\forall I \in L^\infty([0, T], L^2(\Omega)) \text{ with } \text{supp}(I) \subseteq \Omega_{con} \times [0, T] \text{ and } |I(x, t)| \leq R \text{ } (\forall) (x, t) \in \Omega_T.$$

To (5.42), we may apply a Lebesgue point argument analogous to [KUNISCH/WAGNER 12], p. 1541, Proof of Corollary 3.6., in order to get

$$\left(\mu \cdot Q\hat{I}(x_0, t_0) - Q P_2(x_0, t_0) \right) \cdot (\eta_0 - \hat{I}(x_0, t_0)) \geq 0 \text{ } \forall \eta_0 \in [-R, R] \text{ } (\forall) (x_0, t_0) \in \Omega_{con} \times [0, T], \quad (5.43)$$

and this implies the conditions (5.4) and (5.5). ■

Proof of Corollary 5.4. This is implied by (5.5) since $Q P_2 |_{\Omega_{con}} \in L^2[(0, T), W^{1,2}(\Omega_{con})]$ together with $P_2 \in L^2[(0, T), W^{1,2}(\Omega)]$. ■

Proof of Corollary 5.5. We can follow the proof of Theorem 5.2. where only in (5.36), (5.38) and (5.39) the minorization by 0 must be deleted. ■

Acknowledgement.

The present work has been supported within the Special Research Unit 32 “Mathematical Optimization and Applications in Biomedical Sciences” (Graz) by the Austrian Science Fund and the project “Relaxation theorems and necessary optimality conditions for semiconvex multidimensional control problems” (Leipzig) by the German Research Council.

References.

1. [AINSEBA/BENDAHDANE/RUIZ-BAIER 12] Ainseba, B.; Bendahmane, M.; Ruiz-Baier, R.: *Analysis of an optimal control problem for the tridomain model in cardiac electrophysiology*. J. Math. Anal. Appl. **388** (2012), 231 – 247
2. [ALIEV/PANFILOV 96] Aliev, R. R.; Panfilov, A. V.: *A simple two-variable model of cardiac excitation*. Chaos, Solitons & Fractals **7** (1996), 293 – 301
3. [BERGER 77] Berger, M. S.: *Nonlinearity and Functional Analysis*. Academic Press; New York - San Francisco - London 1977
4. [BOURGAULT/COUDIÈRE/PIERRE 09] Bourgault, Y.; Coudière, Y.; Pierre, C.: *Existence and uniqueness of the solution for the bidomain model used in cardiac electrophysiology*. Nonlinear Analysis: Real World Appl. **10** (2009), 458 – 482
5. [BRANDAÕ/FERNÁNDEZ-CARA/MAGALHÃES/ROJAS-MEDAR 08] Brandaõ, A. J. V.; Fernández-Cara, E.; Magalhães, P. M. D.; Rojas-Medar, M. A.: *Theoretical analysis and control results for the FitzHugh-Nagumo equation*. Electronic J. of Differential Equations, Vol. 2008, No. 164, 1 – 20 (electronically published)
6. [DACOROGNA 08] Dacorogna, B.: *Direct Methods in the Calculus of Variations*. Springer; New York etc. 2008, 2nd ed.
7. [EVANS 98] Evans, L. C.: *Partial Differential Equations*. American Mathematical Society; Providence 1998
8. [FITZHUGH 61] FitzHugh, R.: *Impulses and physiological states in theoretical models of nerve membrane*. Biophysical J. **1** (1961), 445 – 466
9. [ITO/KUNISCH 08] Ito, K.; Kunisch, K.: *Lagrange Multiplier Approach to Variational Problems and Applications*. SIAM; Philadelphia 2008
10. [KUNISCH/NAGAIHAH/WAGNER 12] Kunisch, K.; Nagaiah, C.; Wagner, M.: *A parallel Newton-Krylov method for optimal control of the monodomain model in cardiac electrophysiology*. Computing and Visualization in Science **14** (2011) [2012], 257 – 269
11. [KUNISCH/WAGNER 12] Kunisch, K.; Wagner, M.: *Optimal control of the bidomain system (I): The monodomain approximation with the Rogers-McCulloch model*. Nonlinear Analysis: Real World Applications **13** (2012), 1525 – 1550
12. [KUNISCH/WAGNER 11] Kunisch, K.; Wagner, M.: *Optimal control of the bidomain system (II): Uniqueness and regularity theorems*. University of Graz, Institute for Mathematics and Scientific Computing, SFB-Report No. 2011–008 (to appear: Ann. Mat. Pura Appl.)
13. [MUZDEKA/BARBIERI 05] Muzdeka, S.; Barbieri, E.: *Control theory inspired considerations for the mathematical model defibrillation*. In: Proceedings of the 44th IEEE Conference on Decision and Control, 2005 and 2005 European Control Conference, 7416 – 7421
14. [NAGAIHAH/KUNISCH 11] Nagaiah, C.; Kunisch, K.: *Higher order optimization and adaptive numerical solution for optimal control of monodomain equations in cardiac electrophysiology*. Appl. Num. Math. **61** (2011), 53 – 65

-
15. [NAGAIAH/KUNISCH/PLANK 11A] Nagaiah, C.; Kunisch, K.; Plank, G.: *Numerical solution for optimal control of the reaction-diffusion equations in cardiac electrophysiology*. *Comput. Optim. Appl.* **49** (2011), 149 – 178
 16. [NAGAIAH/KUNISCH/PLANK 11B] Nagaiah, C.; Kunisch, K.; Plank, G.: *Optimal control approach to termination of re-entry waves in cardiac electrophysiology*. University of Graz, Institute for Mathematics and Scientific Computing, SFB-Report No. 2011–020 (to appear: *J. Math. Biology*, doi: 10.1007/s00285-012-0557-2)
 17. [NAGUMO/ARIMOTO/YOSHIZAWA 62] Nagumo, J.; Arimoto, S.; Yoshizawa, S.: *An active pulse transmission line simulating nerve axon*. *Proceedings of the Institute of Radio Engineers* **50** (1962), 2061 – 2070
 18. [ROGERS/MCCULLOCH 94] Rogers, J. M.; McCulloch, A. D.: *A collocation-Galerkin finite element model of cardiac action potential propagation*. *IEEE Transactions of Biomedical Engineering* **41** (1994), 743 – 757
 19. [ROLEWICZ 76] Rolewicz, S.: *Funktionalanalysis und Steuerungstheorie*. Springer; Berlin - Heidelberg - New York 1976
 20. [SUNDNES/LINES/CAI/NIELSEN/MARDAL/TVEITO 06] Sundnes, J.; Lines, G. T.; Cai, X.; Nielsen, B. F.; Mardal, K.-A.; Tveito, A.: *Computing the Electrical Activity in the Heart*. Springer; Berlin etc. 2006
 21. [TUNG 78] Tung, L.: *A Bi-Domain Model for Describing Ischemic Myocardial D-C Potentials*. PhD thesis. Massachusetts Institute of Technology 1978
 22. [VENERONI 09] Veneroni, M.: *Reaction-diffusion systems for the macroscopic bidomain model of the cardiac electric field*. *Nonlinear Analysis: Real World Applications* **10** (2009), 849 – 868
 23. [WARGA 72] Warga, J.: *Optimal Control of Differential and Functional Equations*. Academic Press; New York - London 1972
 24. [YOSIDA 95] Yosida, K.: *Functional Analysis*. Springer; Berlin etc. 1995 (reprint of the 6th ed. from 1980)

Last modification: 27.05.2013

Authors' addresses / e-mail. *Karl Kunisch:* University of Graz, Institute for Mathematics and Scientific Computing, Heinrichstraße 36, A-8010 Graz, Austria. e-mail: karl.kunisch@uni-graz.at

Marcus Wagner: University of Leipzig, Department of Mathematics, P. O. B. 10 09 20, D-04009 Leipzig, Germany. Homepage / e-mail: www.thecityto come.de / marcus.wagner@math.uni-leipzig.de